

## On controllability for a class of second-order differential inclusions

AURELIAN CERNEA

### ABSTRACT.

By using a suitable fixed point theorem a sufficient condition for controllability is obtained for a Sturm-Liouville type differential inclusion in the case when the right hand side has convex values.

### REFERENCES

- [1] Benchohra, M. and Ntouyas, S. K., *Controllability for functional differential and integrodifferential inclusions in Banach spaces*, J. Optim. Theory Appl. **113** (2002), 449–472
- [2] Castaing, C. and Valadier, M., *Convex Analysis and Measurable Multifunctions*, Lectures Notes in Mathematics **580**, Springer, Berlin, 1977
- [3] Cernea, A. *Continuous version of Filippov theorem for a Sturm-Liouville type differential inclusion*, Electronic J. Diff. Eq. **2008** (2008), No. 53, 1–7
- [4] Cernea, A. *Sturm-Liouville type differential inclusions in non separable Banach spaces*, Math. Reports, **10** (60) (2008), 205–211
- [5] Cernea, A. *A Filippov type existence theorem for a class of second-order differential inclusions*, J. Inequal. Pure Appl. Math. **9** (2008), No.2, 1–6
- [6] Chang, Y. K. and Li, W. T., *Existence results for second order impulsive functional differential inclusions*, J. Math. Anal. Appl. **301** (2005), 477–490
- [7] Lasota A. and Opial Z., *An application of the Kakutani-Ky-Fan theorem in the theory of ordinary differential equations*, Bull. Acad. Polon. Sci. Math., Astronom. Physiques, **13** (1965), 781–786
- [8] Liu, Y., Wu, J. and Li, Z., *Impulsive boundary value problems for Sturm-Liouville type differential inclusions*, J. Sys. Sci. Complexity **20** (2007), 370–380
- [9] Martelli, M., *A Rothe's type theorem for noncompact acyclic-valued map*, Boll. Unione Matemat. Italiana **11** (1975), 70–76

FACULTY OF MATHEMATICS AND INFORMATICS  
UNIVERSITY OF BUCHAREST  
ACADEMIEI 14, 010014 BUCHAREST, ROMANIA  
E-mail address: acernea@fmi.unibuc.ro