

Remarks on a new metric in the unity disc of the complex plane

LAURIAN-IOAN PIȘCORAN and CĂTĂLIN BARBU

ABSTRACT.

The curvature K of a surface depends only to the surface metric and so is an intrinsic invariant. In this paper we will study a new metric in the unity complex disc, which is connected to the well-known Poincaré metric used in hyperbolic geometry.

REFERENCES

- [1] Bercu, G., Matsuyama, Y. and Postolache M., *Hessian Metrics and Ricci Solitons*, Fair Parteners Publishers, București, 2011
- [2] Krantz, S. G., *Geometric Function Theory Explorations in Complex Analysis*, Birkhauser, Boston, 2006
- [3] Isaev, A. V. and Krantz, S. G., *Invariant Distances and Metrics in Complex Analysis*, Notices of the AMS, **47** (2000), No. 5, 546–553
- [4] Pișcoran, L. I. and Barbu, C. I., *A new hyperbolic metric*, Differential Geometry-Dynamical Systems, **15** (2013), 71–78

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE
FACULTY OF SCIENCE NORTH UNIVERSITY CENTER AT BAIA MARE
TECHNICAL UNIVERSITY OF CLUJ NAPOCA
VICTORIEI 76, 430122 BAIA MARE, ROMANIA
E-mail address: plaurian@yahoo.com

"VASILE ALECSANDRI" NATIONAL COLLEGE
VASILE ALECSANDRI 37, 600011 BACĂU, ROMANIA
E-mail address: kafka_mate@yahoo.com

Received: 30.04.2013; In revised form: 16.10.2013; Accepted: 15.01.2014

2010 *Mathematics Subject Classification.* 30F45, 20N99, 51B10, 51M10.

Key words and phrases. *Hyperbolic geometry, complex disk, Poincaré metric.*

Corresponding author: Laurian-Ioan Pișcoran; plaurian@yahoo.com