Fixed point theorems for multivalued contractions in Kasahara spaces

ALEXANDRU-DARIUS FILIP

Abstract.

In this paper we present some fixed point results for multivalued α -contractions. Our results are obtained in a more general setting, the so called Kasahara space. Some of them are generalizations of Maia's type fixed point result for multivalued α -contractions. As application, a fixed point theorem for integral inclusions is given.

REFERENCES

- [1] Aubin, J. P. and Frankowska, H., Set-Valued Analysis, Birkhäuser, Boston, 1990
- [2] Fréchet, M., Les espaces abstraits, Gauthier-Villars, Paris, 1928
- [3] Kasahara, S., Common fixed points for multivalued mappings in L-spaces, Math. Sem. Notes, 4 (1976), 181–193
- [4] Kasahara, S., On some generalizations of the Banach contraction theorem, Publ. RIMS, Kyoto Univ., 12 (1976), 427–437
- [5] Markin, J. T., A fixed point theorem for set-valued mappings, Bull. Amer. Math. Soc., 74 (1968), 639-640
- [6] Nadler, S. B., Multi-valued contraction mappings, Pacific J. Math., 30 (1969), 282-291
- [7] Petruşel, A. and Rus, I. A., Fixed point theory for multivalued operators on a set with two metrics, Fixed Point Theory, 8 (2007), No. 1, 97–104
- [8] Rus, I. A., Picard operators and applications, Sci. Math. Japonicae, 58 (2003), 191-219
- [9] Rus, I. A., Kasahara spaces, Sci. Math. Jpn., 72 (2010), No. 1, 101–110

DEPARTMENT OF MATHEMATICS BABEŞ-BOLYAI UNIVERSITY KOGĂLNICEANU 1, 400084 CLUJ-NAPOCA, ROMANIA *E-mail address*: darius.filip@econ.ubbcluj.ro

Received: 15.08.2013; In revised form: 28.04.2014; Accepted: 30.04.2014 2010 Mathematics Subject Classification. 47H10, 54H25.

Key words and phrases. Kasahara space, multivalued α -contraction, fixed point, Pompeiu-Hausdorff functional, multivalued weakly Picard operator, sequence of successive approximations, Fredholm-type integral inclusion.