CARPATHIAN J. MATH.
Volume **36** (2020), No. 3,
Pages 373 - 379

Online version at https://www.carpathian.cunbm.utcluj.ro/
Print Edition: ISSN 1584 - 2851; Online Edition: ISSN 1843 - 4401
DOI: https://doi.org/10.37193/CJM.2020.03.05

New univalence criteria for an integral operator with Mocanu's and Şerb's lemma

CAMELIA BĂRBATU and DANIEL BREAZ

ABSTRACT. In this paper we consider a general integral operator for analytic functions in the open unit disk $\mathbb{U}$ and we obtain sufficient conditions for univalence of this integral operator, using Mocanu's and Şerb's Lemma. This integral operator was considered in a recent work Bărbatu, C. and Breaz, D., [*The univalence conditions for a general integral operator*, Acta Univ. Apulensis Math. Inform., **57** (2019), 133–145]. The results derived in this paper are shown to follow upon specializing the parameters involved in our results. Several corollaries of the main results are also considered.

1. INTRODUCTION AND PRELIMINARIES

Let $\mathcal{A}$ denote the class of the functions of the form:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \tag{1.1}$$

which are analytic in the open unit disk

$$\mathbb{U} = \{z \in \mathbb{C} :\mid z \mid < 1\}$$

and satisfy the following usual normalization conditions:

$$f(0) = f^{'}(0) - 1 = 0,$$

$\mathbb{C}$ being the set of complex numbers. We denote by $\mathcal{S}$ the subclass of $\mathcal{A}$ consisting of functions $f \in \mathcal{A}$, which are univalent in $\mathbb{U}$.

We consider the integral operator:

$$\mathcal{M}_n(f_i, g_i)(z) = \left\{ \delta \int_0^z t^{\delta-1} \prod_{i=1}^{n} \left[\left(\frac{f_i(t)}{t} \right)^{\alpha_i - 1} (g_i{}'(t))^{\beta_i} \left(\frac{g_i(t))}{t} \right)^{\gamma_i} \right] \mathrm{dt} \right\}^{\frac{1}{\delta}}, \tag{1.2}$$

where f_i, g_i are analytic in $\mathbb{U}$, and $\alpha_i, \beta_i, \gamma_i \in \mathbb{C}$ for all $i = \overline{1,n}$, $n \in \mathbb{N} \setminus \{0\}$, $\delta \in \mathbb{C}$, with $\mathrm{Re}\delta > 0$.

Remark 1.1. The integral operator $\mathcal{M}_n(f_i, g_i)$ defined by (1.2), introduced by Bărbatu and Breaz in the paper [1], is a general integral operator of Pfaltzgraff, Kim-Merkes and Oversea types which extends also the other operators as follows:

i) For $n = 1$, $\delta = 1$, $\alpha_1 - 1 = \alpha_1$ and $\beta_1 = \gamma_1 = 0$ we obtain the integral operator which was studied by Kim-Merkes [7]

$$\mathcal{F}_\alpha(z) = \int_0^z \left(\frac{f(t)}{t} \right)^{\alpha} \mathrm{dt}.$$

Received: 20.10.2019. In revised form: 04.05.2020. Accepted: 11.05.2020
2010 *Mathematics Subject Classification.* 30C45, 30C75.
Key words and phrases. *Integral operators, analytic and univalent functions, open unit disk, univalence conditions, Mocanu's and Şerb's Lemma, Schwarz Lemma.*
Corresponding author: Camelia Bărbatu; camipode@yahoo.com

ii) For $n = 1$, $\delta = 1$ and $\alpha_1 - 1 = \gamma_1 = 0$ we obtain the integral operator which was studied by Pfaltzgraff [20]

$$\mathcal{G}_\alpha(z) = \int_0^z \left(f'(t)\right)^\alpha \mathrm{dt}.$$

iii) For $\alpha_i - 1 = \alpha_i$ and $\beta_i = \gamma_i = 0$ we obtain the integral operator which was defined and studied by D. Breaz and N. Breaz [2]

$$\mathcal{D}_n(z) = \left[\delta \int_0^z t^{\delta-1} \prod_{i=1}^n \left(\frac{f_i(t)}{t}\right)^{\alpha_i} \mathrm{dt}\right]^{\frac{1}{\delta}},$$

this integral operator is a generalization of the integral operator introduced by Pascu and Pescar [16].

iv) For $\alpha_i - 1 = \gamma_i = 0$ we obtain the integral operator which was defined and studied by D. Breaz, Owa and N. Breaz [5]

$$\mathcal{I}_n(z) = \left[\delta \int_0^z t^{\delta-1} \prod_{i=1}^n \left[f_i'(t)\right]^{\alpha_i} \mathrm{dt}\right]^{\frac{1}{\delta}},$$

this integral operator is a generalization of the integral operator introduced by Pescar and Owa in [19] .

v) For $\alpha_i - 1 = \alpha_i$ and $\gamma_i = 0$ we obtain the integral operator which was defined and studied by Pescar in [17]

$$\mathcal{F}_n(z) = \left[\delta \int_0^z t^{\delta-1} \prod_{i=1}^n \left(\frac{f_i(t)}{t}\right)^{\alpha_i} \left(f_i'(t)\right)^{\beta_i} \mathrm{dt}\right]^{\frac{1}{\delta}},$$

this integral operator is a generalization of the integral operator introduced by Frasin in [6] and by Oversea in [12].

vi) For $\alpha_i - 1 = \alpha_i$ and $\gamma_i = 0$ we obtain the integral operator which was studied by Ularu in [21]

$$\mathcal{I}_n(z) = \left[\delta \int_0^z t^{\delta-1} \prod_{i=1}^n \left(\frac{f_i(t)}{t}\right)^{\alpha_i} \left(g_i'(t)\right)^{\beta_i} \mathrm{dt}\right]^{\frac{1}{\delta}}.$$

Thus, the integral operator $\mathcal{M}_n(f_i, g_i)$, introduced here by the formula (1.2), can be considered as an extension and a generalization of these operators above mentioned.

The following two univalence conditions were derived by Pascu [14] and [15].

Theorem 1.1. (Pascu [14]) *Let $f \in \mathcal{A}$ and $\gamma \in \mathbb{C}$. If $Re\gamma > 0$ and*

$$\frac{1 - |z|^{2Re\gamma}}{Re\gamma} \left|\frac{zf''(z))}{f'(z)}\right| \leq 1,$$

for all $z \in \mathbb{U}$, then the integral operator

$$F_\gamma(z) = \left(\gamma \int_0^z t^{\gamma-1} f'(t) dt\right)^{\frac{1}{\gamma}},$$

is in the class $\mathcal{S}$.

Theorem 1.2. (Pascu [15]) *Let $\delta \in \mathbb{C}$ with $Re\delta > 0$. If $f \in \mathcal{A}$ satisfies*

$$\frac{1 - |z|^{2Re\delta}}{Re\delta} \left|\frac{zf''(z))}{f'(z)}\right| \leq 1,$$

for all $z \in \mathbb{U}$, then, for any complex γ with $Re\gamma \geq Re\delta$, the integral operator

$$F_\gamma(z) = \left(\gamma \int_0^z t^{\gamma-1} f'(t)dt\right)^{\frac{1}{\gamma}},$$

is in the class $\mathcal{S}$.

Mocanu and Şerb, on the other hand, proved the next Theorem in [10].

Theorem 1.3. (Mocanu - Şerb [10]) *Let $M_0 = 1,5936...$ the positive solution of equation*

$$(2 - M)\, e^M = 2. \tag{1.3}$$

If $f \in \mathcal{A}$ and

$$\left|\frac{f''(z))}{f'(z)}\right| \leq M_0,$$

for $z \in \mathbb{U}$, then

$$\left|\frac{zf'(z))}{f(z)} - 1\right| \leq 1, \qquad (z \in \mathbb{U})$$

The bound M_0 is sharp.

Finally, in our present investigation, we shall also need the familiar Schwarz Lemma [8].

Lemma 1.1. (General Schwarz Lemma [8]) *Let f be the function regular in the disk $\mathbb{U}_R = \{z \in \mathbb{C} : |z| < R, R > 0\}$ with $|f(z)| < M$ for a fixed number $\mathrm{M} > 0$ fixed. If $f(z)$ has one zero with multiplicity order bigger than a positive integer m for $z = 0$, then*

$$|f(z)| \leq \frac{M}{R^m} z^m, \quad z \in \mathbb{U}_R.$$

The equality for $z \neq 0$ can hold only if

$$f(z) = e^{i\theta} \frac{M}{R^m} z^m,$$

where θ is constant.

2. The main univalence criterion

In this paper, we obtain new conditions for the univalence of the general integral operator $\mathcal{M}_n(f_i, g_i)$, by applying the improvement of Becker univalence criteria, obtained by Pascu in the paper [14]. Also, a lemma given by Mocanu and Şerb in the paper [10], will be used to get some part of results.

Theorem 2.4. *Let $\gamma, \delta, \alpha_i, \beta_i, \gamma_i$ be complex numbers, $c = Re\gamma > 0$, $i = \overline{1,n}$, M_0 the positive solution of the equation (1.3), $M_0 = 1,5936...$ and $f_i, g_i \in \mathcal{A}$. If*

$$\left|\frac{f_i''(z)}{f_i'(z)}\right| \leq M_0, \quad \left|\frac{g_i''(z)}{g_i'(z)}\right| \leq M_0, \tag{2.4}$$

for all $z \in \mathbb{U}$, $i = \overline{1,n}$ and

$$\frac{1}{c}\sum_{i=1}^{n} |\alpha_i - 1| + \frac{2M_0}{(2c+1)^{\frac{2c+1}{2c}}} \sum_{i=1}^{n} |\beta_i| + \frac{1}{c}\sum_{i=1}^{n} |\gamma_i| \leq 1. \tag{2.5}$$

Then, for all δ complex numbers, $Re\delta \geq Re\gamma$, the integral operator $\mathcal{M}_n(f_i, g_i)$, given by (1.2) is in the class $\mathcal{S}$.

Proof. Let us define the function

$$M_n(z) = \int_0^z \prod_{i=1}^n \left(\frac{f_i(t)}{t}\right)^{\alpha_i - 1} (g_i'(t))^{\beta_i} \left(\frac{g_i(t)}{t}\right)^{\gamma_i} \mathrm{dt},$$

for $f_i, g_i \in \mathcal{A}$, $i = \overline{1,n}$.

The function M_n is regular in $\mathbb{U}$ and satisfy the following usual normalization conditions $M_n(0) = M_n^{'}(0) - 1 = 0$.

After some calculus we have

$$\frac{zM_n''(z)}{M_n'(z)} = \sum_{i=1}^n \left[(\alpha_i - 1)\left(\frac{zf_i'(z)}{f_i(z)} - 1\right) + \beta_i \frac{zg_i''(z)}{g_i'(z)} + \gamma_i \left(\frac{zg_i'(z)}{g_i(z)} - 1\right)\right],$$

for all $z \in \mathbb{U}$.

Applying the module and multiplying both members with $\frac{1-|z|^{2c}}{c}$, we obtain

$$\frac{1-|z|^{2c}}{c}\left|\frac{zM_n''(z)}{M_n'(z)}\right| \leq$$

$$\leq \frac{1-|z|^{2c}}{c} \sum_{i=1}^n \left[|\alpha_i - 1|\left|\frac{zf_i'(z)}{f_i(z)} - 1\right| + |\beta_i|\,|z|\left|\frac{g_i''(z)}{g_i'(z)}\right| + |\gamma_i|\left|\frac{zg_i'(z)}{g_i(z)} - 1\right|\right], \tag{2.6}$$

for all $z \in \mathbb{U}$.

Using (2.4) and Lemma Mocanu and Șerb, from (2.6) we get

$$\left|\frac{zf_i'(z)}{f_i(z)} - 1\right| < 1, \quad \left|\frac{zg_i'(z)}{g_i(z)} - 1\right| < 1,$$

for all $z \in \mathbb{U}$, $i = \overline{1,n}$ and hence, we have

$$\frac{1-|z|^{2c}}{c}\left|\frac{zM_n''(z)}{M_n'(z)}\right| \leq$$

$$\leq \frac{1-|z|^{2c}}{c} \sum_{i=1}^n |\alpha_i - 1| + \frac{1-|z|^{2c}}{c}\,|z|\,M_0 \sum_{i=1}^n |\beta_i| + \frac{1-|z|^{2c}}{c} \sum_{i=1}^n |\gamma_i|, \tag{2.7}$$

for all $z \in \mathbb{U}$.

Since

$$\max_{|z|\leq 1} \frac{\left(1-|z|^{2c}\right)|z|}{c} = \frac{2}{(2c+1)^{\frac{2c+1}{2c}}}, \tag{2.8}$$

from (2.7) and (2.8) we obtain

$$\frac{1-|z|^{2c}}{c}\left|\frac{zM_n''(z)}{M_n'(z)}\right| \leq \frac{1}{c}\sum_{i=1}^n |\alpha_i - 1| + \frac{2M_0}{(2c+1)^{\frac{2c+1}{2c}}} \sum_{i=1}^n |\beta_i| + \frac{1}{c}\sum_{i=1}^n |\gamma_i|, \tag{2.9}$$

for all $z \in \mathbb{U}$.

Using (2.5), from (2.9) we have

$$\frac{1-|z|^{2c}}{c}\left|\frac{zM_n''(z)}{M_n'(z)}\right| \leq 1, \tag{2.10}$$

for all $z \in \mathbb{U}$.

By Theorem 1.2 it results that $\mathcal{M}_n(f_i, g_i) \in \mathcal{S}$.

□

3. Corollaries and consequences

First of all, upon setting $\delta = 1$ in Theorem 2.4, we obtain the following corollary:

Corollary 3.1. *Let $\gamma, \alpha_i, \beta_i, \gamma_i$ be complex numbers, $0 < Re\gamma \leq 1$, $c = Re\gamma$, M_0 the positive solution of the equation (1.3), $M_0 = 1,5936...$ and $f_i, g_i \in \mathcal{A}$. If*

$$\left|\frac{f_i''(z)}{f_i'(z)}\right| \leq M_0, \quad \left|\frac{g_i''(z)}{g_i'(z)}\right| \leq M_0,$$

for all $z \in \mathbb{U}$, $i = \overline{1,n}$ and

$$\frac{1}{c}\sum_{i=1}^{n}|\alpha_i - 1| + \frac{2M_0}{(2c+1)^{\frac{2c+1}{2c}}}\sum_{i=1}^{n}|\beta_i| + \frac{1}{c}\sum_{i=1}^{n}|\gamma_i| \leq 1$$

then, the integral operator $\mathcal{M}_n^*$, defined by

$$\mathcal{M}_n^*(z) = \int_0^z \coprod_{i=1}^{n}\left[\left(\frac{f_i(t)}{t}\right)^{\alpha_i - 1}(g_i'(t))^{\beta_i}\left(\frac{g_i(t))}{t}\right)^{\gamma_i}\right]dt \tag{3.11}$$

is in the class $\mathcal{S}$.

Letting $\delta = 1$ and $\gamma_i = 0$ in Theorem 2.4, we have:

Corollary 3.2. *Let $\gamma, \alpha_i, \beta_i$ be complex numbers, $0 < Re\gamma \leq 1$, $c = Re\gamma$, M_0 the positive solution of the equation (1.3), $M_0 = 1,5936...$ and $f_i, g_i \in \mathcal{A}$. If*

$$\left|\frac{f_i''(z)}{f_i'(z)}\right| \leq M_0, \quad \left|\frac{g_i''(z)}{g_i'(z)}\right| \leq M_0,$$

for all $z \in \mathbb{U}$, $i = \overline{1,n}$ and

$$\frac{1}{c}\sum_{i=1}^{n}|\alpha_i - 1| + \frac{2M_0}{(2c+1)^{\frac{2c+1}{2c}}}\sum_{i=1}^{n}|\beta_i| \leq 1$$

then, the integral operator $\mathcal{F}_n$ defined by

$$\mathcal{F}_n(z) = \int_0^z \coprod_{i=1}^{n}\left[\left(\frac{f_i(t)}{t}\right)^{\alpha_i - 1}(g_i'(t))^{\beta_i}\right]dt \tag{3.12}$$

is in the class $\mathcal{S}$.

Remark 3.2. The integral operator from (3.12) is a known result, proven in [21].

Letting $\delta = 1$ and $\beta_i = 0$ in Theorem 3.1, we have

Corollary 3.3. *Let $\gamma, \alpha_i, \gamma_i$be complex numbers, $0 < Re\gamma \leq 1$, $c = Re\gamma$, M_0 the positive solution of the equation (1.3), $M_0 = 1,5936...$ and $f_i, g_i \in \mathcal{A}$, $i = \overline{1,n}$. If*

$$\left|\frac{f_i''(z)}{f_i'(z)}\right| \leq M_0, \quad \left|\frac{g_i''(z)}{g_i'(z)}\right| \leq M_0,$$

for all $z \in \mathbb{U}$, $i = \overline{1,n}$ and

$$\frac{1}{c}\sum_{i=1}^{n}|\alpha_i - 1| + \frac{1}{c}\sum_{i=1}^{n}|\gamma_i| \leq 1$$

then, the integral operator $\mathcal{G}_n$ defined by

$$\mathcal{G}_n(z) = \int_0^z \coprod_{i=1}^{n}\left[\left(\frac{f_i(t)}{t}\right)^{\alpha_i - 1}\left(\frac{g_i(t))}{t}\right)^{\gamma_i}\right]dt \tag{3.13}$$

is in the class $\mathcal{S}$.

Remark 3.3. The integral operator from (3.13) is another known result proven in [11].

Letting $\delta = 1$ and $\alpha_i = 1$ in Theorem 2.4, we obtain the next corollary:

Corollary 3.4. *Let $\gamma, \beta_i, \gamma_i$ be complex numbers, $0 < Re\gamma \leq 1$, $c = Re\gamma$, M_0 the positive solution of the equation (1.3), $M_0 = 1,5936...$ and $g_i \in \mathcal{A}$. If*

$$\left|\frac{g_i''(z)}{g_i'(z)}\right| \leq M_0,$$

for all $z \in \mathbb{U}$, $i = \overline{1,n}$ and

$$\frac{2M_0}{(2c+1)^{\frac{2c+1}{2c}}} \sum_{i=1}^{n} |\beta_i| + \frac{1}{c} \sum_{i=1}^{n} |\gamma_i| \leq 1$$

then, the integral operator $\mathcal{I}_n$ defined by

$$\mathcal{I}_n(z) = \int_0^z \prod_{i=1}^{n} \left[(g_i'(t))^{\beta_i} \left(\frac{g_i(t))}{t} \right)^{\gamma_i} \right] dt \tag{3.14}$$

is in the class $\mathcal{S}$.

Remark 3.4. The integral operator given by (3.14) was proved in [17].

Letting $n = 1$, $\delta = \gamma = \alpha$ and $\alpha_i - 1 = \beta_i = \gamma_i$ in Theorem 2.4, we obtain:

Corollary 3.5. *Let α be complex numbers, $a = Re\alpha > 0$, M_0 the positive solution of the equation (1.3), $M_0 = 1,5936...$ and $f, g \in \mathcal{A}$. If*

$$\left|\frac{f''(z)}{f'(z)}\right| \leq M_0, \quad \left|\frac{g''(z)}{g'(z)}\right| \leq M_0,$$

for all $z \in \mathbb{U}$, and

$$2\left(\alpha - 1\right)\left(\frac{1}{a} + \frac{M_0}{(2a+1)^{\frac{2a+1}{2a}}}\right) \leq 1,$$

then the integral operator $\mathcal{M}$ defined by

$$\mathcal{M}(z) = \left\{ \alpha \int_0^z \left[f(t) g'(t) \frac{g(t)}{t} \right]^{\alpha - 1} dt \right\}^{\frac{1}{\alpha}}, \tag{3.15}$$

is in the class $\mathcal{S}$.

4. Conclusions and further study

The future research will study similar properties using other classes of analytical functions for the integral operators which were defined in this paper, further generalizing the integral operators and getting results related to convexity, starlikeness, univalence conditions, etc.

References

[1] Bărbatu, C. and Breaz, D., *The univalence conditions for a general integral operator*, Acta Univ. Apulensis Math. Inform., **57** (2019), 133–145

[2] Breaz, D. and Breaz, N., *Two Integral Operators*, Studia Univ. Babes-Bolyai, Math., **47** (2002), No. 3, 13–21

[3] Breaz, D. and Breaz, N., *Univalence conditions for certain integral operators*, Studia Univ. Babeş-Bolyai Math., **2** (2002), 9–17

[4] Breaz, D., Breaz, N. and Srivastava H. M., *An extension of the univalent condition for a family of integral operators*, Appl. Math. Lett., **22** (2009), No. 3, 41–44

[5] Breaz, D., Owa, S. and Breaz, N., *A new integral univalent operator*, Acta Univ. Apulensis Math. Inform., **16** (2008), 11–16

[6] Frasin, B. A., *Order of convexity and univalence of general integral operator*, J. Franklin Inst., **348** (2011), 1013–1019
[7] Kim, Y. J. and Merkes, E. P., *On an integral of powers of a spirallike function*, Kyungpook Math. J., **12** (1972), 249–253
[8] Mayer, O., *The Functions Theory of the One Variable Complex* , Acad. Ed. Bucureşti, 1981, 101–117
[9] Mocanu, P. T., Bulboaca, T. and Salagean, G. S., *Teoria geometrică a funcţiilor univalente*, Casa Cărţii de Ştiinţă, Cluj Napoca (1999), 77–81
[10] Mocanu, P. T. and Şerb I., *A sharp simple criterion for a subclass of starlike functions*, Complex variables, **32** (1997), 161–168
[11] Oprea, A. and Breaz, D., *Univalence conditions for a general operator*, An. Șt. Univ. Ovidius Constanța, **23** (2015), No. 1, 213–224
[12] Oversea, H., *Integral operators of Bazilvic type*, Bull. Math. Bucureşti, **37** (1993), 115–125
[13] Ozaki, S. and Nunokawa, M., *The Schwarzian derivative and univalent functions*, Proceed. American Math. Soc. Math., **33** (1972), 392–394
[14] Pascu, N. N., *An a univalence criterion II*, Itinerant Seminar on Functional Equations, Approximation and Convexity, Cluj Napoca 1985, 153–154
[15] Pascu, N. N., An improvement of Becker's univalence criterion, *Proceedings of the Commemorative Session Simion Stoilow*, Braşov, Preprint (1987), 43–48
[16] Pascu, N. N. and Pescar, V., *On the integral operators Kim-Merkes and Pfaltzgraff*, Mathematica, Univ. Babeş-Bolyai, Cluj-Napoca, **32** (55) (1990), No. 2, 185–192
[17] Pescar V., *New univalence criteria for some integral operators*. Studia Univ. Babes-Bolyai Math. **59** (2014), No. 2, 185-192.
[18] Pescar, V. and Breaz, N., *Mocanu and Şerb type univalence criteria for some general integral operators*, Acta Univ. Apulensis Math. Inform., **44** (2015), 1–8
[19] Pescar, V. and Owa, S., *Univalence of certain integral operators*, Int. J. Math. Math. Sci., **23** (2000), 697–701
[20] Pfaltzgraff, J. A., *Univalence of the integral* $f'(z)^{\lambda}$, Bul. London Math. Soc., **7** (1975), No. 3, 254–256
[21] Ularu, N., *Convexity properties for an integral operator*, Acta Univ. Apulensis Math. Inform., **27** (2011), 115–120

"Babeş-Bolyai" University
Faculty of Mathematics and Computer Sciences
Kogălniceanu 1, 400084 Cluj-Napoca, Romania

Department of Mathematics
"1 Decembrie" University
Gabriel Bethlen 5, 510009 Alba-Iulia, Romania
Email address: camipode@yahoo.com
Email address: dbreaz@uab.ro