

*Dedicated to Professor Yeol Je Cho on the occasion of his retirement*

## Existence and stability for a generalized differential mixed quasi-variational inequality

WEI LI<sup>1</sup>, YI-BIN XIAO<sup>2</sup>, XING WANG<sup>3</sup> and JUN FENG<sup>4</sup>

**ABSTRACT.** In the present paper, we investigate a generalized differential mixed quasi-variational inequality consisting of a system of an ordinary differential equation and a generalized mixed quasi-variational inequality. By using an important result concerning the measurable selection, we prove the existence of Carathéodory weak solution to the generalized differential mixed quasi-variational inequality. Then, with the existence result, we establish two stability results for the generalized differential mixed quasi-variational inequality under different conditions, i.e., upper semicontinuity and lower semicontinuity of the Carathéodory weak solution with respect to the parameter, which is a perturbation of some mappings in the generalized mixed quasi-variational inequality.

### 1. INTRODUCTION

It is well known that variational inequality (VI), which has various kinds of generalizations such as quasi-variational inequality, mixed variational inequality and vector variational inequality etc., has been widely studied and applied into many research fields such as economics, optimization, mechanics and transportation etc. (see [1, 5, 7, 10, 15, 19, 20, 25, 26, 27, 28, 29, 30]). When an ordinary differential equation is involved, the system of variational inequality and ordinary differential equation is called differential variational inequality (DVI), which is introduced and studied by Pang and Stewart [16] in 2008. Recently, DVIs have attracted much attention of researchers in different research fields and many theoretical results, numerical algorithms and applications of DVIs have been studied by many authors. For the literature on the research of DVIs, we refer the reader to [14, 17, 18, 24] and the references therein.

The stability analysis of a VI or a DVI with perturbed data is concerned with the upper and lower semicontinuity, continuity, Lipschitz continuity or some kind of differentiability of its solution set. It can help in identifying sensitive parameters that should be obtained with relatively high accuracy, predicting the future changes of the equilibria as a result of the changes in the governing system, providing useful information for designing or planning various equilibrium systems. Consequently, the stability analysis of VIs and DVIs attracted the attention of many researchers in very early time. In 1986, Tobin studied the stability analysis for a VI when both the variational inequality function and the feasible region are perturbed in [21]. Khanh and Luu[6] studied the lower semicontinuity and upper semicontinuity of the solution sets and approximate solution sets of parametric multi-valued quasi-variational inequalities in topological vector spaces. Recently, Wang et al.[22] studied some upper semicontinuity and continuity results concerned with the

---

Received: 30.09.2017. In revised form: 10.06.2018. Accepted: 15.07.2018

2010 *Mathematics Subject Classification.* 49J40, 35B35, 35D30.

*Key words and phrases.* generalized differential mixed quasi-variational inequalities, Carathéodory weak solutions, upper semicontinuity, lower semicontinuity, stability.

Corresponding author: Xing Wang; wangxing0793@163.com

Carathéodory weak solution set mapping for the differential set-valued mixed variational inequality. For more related research works, we can refer to [4, 8, 23] and the references therein.

Inspired by the above research on stability analysis of VI and DVI, in this paper, we study the existence and stability of solution to the following generalized differential mixed quasi-variational inequality (GDMQVI) in finite dimensional space:

Find  $(x(t), u(t)) : [0, T] \rightarrow R^n \times R^n$  such that

$$(1.1) \quad \begin{cases} \dot{x}(t) = f(t, x(t)) + B(t, x(t))u(t), \\ \langle y - h(u), G(t, x) + F(u) \rangle + p\varphi(y) - p\varphi(h(u)) \geq 0, & \forall y \in K(u), \\ h(u) \in K(u), \\ x(0) = x_0. \end{cases}$$

where  $\dot{x}(t) = \frac{dx}{dt}$  stands for the derivative of function  $x$  with respect to time variable  $t$ ,  $(f, B, G) : \Omega \rightarrow R^m \times R^{m \times n} \times R^n$  are given functions with  $\Omega = [0, T] \times R^n$  while  $(h, F)$  are functions from  $R^n$  to  $R^n$ ,  $\varphi : R^n \rightarrow (-\infty, +\infty]$  is a functional,  $p$  is a positive real number, and  $K : R^n \rightrightarrows R^n$  is a set-valued mapping such that  $K(u) \subset R^n$  is a closed convex subset for each  $u \in R^n$ .

The following of this paper is as follows. First, in Section 2, we present some preliminaries. Then we give an existence result of Carathéodory weak solutions to the GDMQVI (1.1) in Section 3. At last, in Sect. 4, the upper semicontinuity and lower semicontinuity of Carathéodory weak solution set with respect to the perturbed data in the generalized mixed quasi-variational inequality is established.

## 2. PRELIMINARES

In this section, we introduce some basic notations and preliminary results, and present some definitions and hypotheses for the GDMQVI (1.1).

**Definition 2.1.** (see [12]) Let  $H$  be a real Hilbert space, and  $g, A : H \rightarrow H$  be two single-valued mappings.

(i)  $A$  is said to be  $\lambda$ -strongly monotone on  $H$  if there exists a constant  $\lambda$  such that

$$\langle Ax - Ay, x - y \rangle \geq \lambda \|x - y\|^2, \quad \forall x, y \in H;$$

(ii)  $(A, g)$  is said to be a  $\mu$ -strongly monotone couple on  $H$  if there exists a constant  $\mu > 0$  such that

$$\langle Ax - Ay, g(x) - g(y) \rangle \geq \mu \|x - y\|^2, \quad \forall x, y \in H.$$

**Definition 2.2.** (see[2]) Let  $X, Y$  be two metric spaces. A set-valued mapping  $F : X \rightrightarrows Y$  is called to be

- (i) upper semicontinuous at  $x_0 \in X$  iff for any neighborhood  $N$  of  $F(x_0)$ , there exists an  $\eta > 0$ , such that  $F(x) \subset N$  for every  $x \in B(x_0, \eta)$ ;
- (ii) lower semicontinuous at  $x_0 \in X$  iff for any  $y \in F(x_0)$  and for any sequence  $x_n \in X$  converging to  $x_0$ , there exists a sequence of elements  $y_n \in F(x_n)$  converging to  $y$ .

**Definition 2.3.** (see [12]) Let  $H$  be a real Hilbert space and  $K : H \rightrightarrows H$  be a set-valued mapping such that  $K(u)$  is a closed and convex subset of  $H$  for any  $u \in H$ . The generalized  $f$ -projection of  $z \in H$  on the set  $K(u)$  is defined by

$$P_{K(u)}z = arg \inf_{\xi \in K(u)} (\|z\|^2 - 2\langle z, \xi \rangle + \|\xi\|^2 + 2p\varphi(\xi)), \quad \forall z \in H.$$

**Definition 2.4.** A pair  $(x(t), u(t))$  on  $[0, T]$  is called a Carathéodory weak solution of the GDMQVI (1.1) if

- (1)  $h(u) \in K(u)$ ;
- (2)  $x(t)$  is absolutely continuous on  $[0, T]$  and satisfies the differential equation in the GDMQVI (1.1) for almost all  $t \in [0, T]$ ;
- (3)  $u \in L^2[0, T]$  satisfies the generalized mixed quasi-variational inequality in the GDMQVI (1.1) for every  $t \in [0, T]$ , where  $L^2[0, T]$  denotes the set of all measurable functions  $u : [0, T] \rightarrow R^n$  satisfying  $\int_0^T \|u(t)\|^2 dt < +\infty$ .

**Lemma 2.1.** ([13]). Let  $\varphi : R^n \rightarrow (-\infty, +\infty]$  be a proper lower semicontinuous convex functional. Suppose that the function  $\varphi(u(\cdot))$  is integrable on  $[0, T]$  for every  $u \in L^2[0, T]$ . Then

$$\phi(u) = \int_0^T \varphi(u(t)) dt, \quad u \in L^2[0, T]$$

is a proper lower semicontinuous convex functional.

**Lemma 2.2.** ([16]) Let  $\mathbb{F} : \Omega \rightrightarrows R^m$  be an upper semicontinuous set-valued mapping with nonempty closed convex values. Suppose that there exists a scalar  $\rho^{\mathbb{F}} > 0$  satisfying

$$(2.2) \quad \sup\{\|y\| : y \in \mathbb{F}(t, x)\} \leq \rho^{\mathbb{F}}(1 + \|x\|), \quad \forall (t, x) \in \Omega.$$

For every  $x^0 \in R^n$ , differential inclusion (DI)  $\dot{x} \in \mathbb{F}(t, x), x(0) = x^0$  has a weak solution in the sense of Carathéodory.

**Lemma 2.3.** ([16]) Let  $H : \Omega \times R^m \rightarrow R^m$  be a continuous function and  $U : \Omega \rightrightarrows R^n$  be a closed set-valued mapping such that for some constant  $\eta_U > 0$ ,

$$\sup_{u \in U(t, x)} \|u\| \leq \eta_U(1 + \|x\|), \quad \forall (t, x) \in \Omega.$$

Let  $v : [0, T] \rightarrow R^m$  be a measurable function and  $x : [0, T] \rightarrow R^m$  be a continuous function satisfying  $v(t) \in H(t, x(t), U(t, x(t)))$  for almost all  $t \in [0, T]$ . There exists a measurable function  $u : [0, T] \rightarrow R^n$  such that  $u(t) \in U(t, x(t))$  and  $v(t) = H(t, x(t), u(t))$  for almost all  $t \in [0, T]$ .

At the end of this section, we present the following two hypotheses (A) and (B), which hold for some functions in GDMQVI (1.1) in the rest of this paper.

- (A) Suppose that the functions  $f, B$  and  $G$  are Lipschitz continuous functions on  $\Omega$  with Lipschitz constants  $L_f > 0, L_B > 0$  and  $L_G > 0$ , respectively.
- (B) Suppose that the functions  $f$  and  $B$  are bounded on  $\Omega$  with constants  $\sigma_f$  and  $\sigma_B$ . i.e.,  $\sigma_B \equiv \sup_{(t, x) \in \Omega} \|B(t, x)\| < \infty$  and  $\sigma_f \equiv \sup_{(t, x) \in \Omega} \|f(t, x)\| < \infty$ .

### 3. EXISTENCE OF SOLUTIONS TO THE GDMQVI

In this section, we prove the existence of Carathéodory weak solutions to the GDMQVI (1.1) by using Lemma 2.2 and Lemma 2.3. For this purpose, we define a set-valued mapping  $\mathbb{F}(t, x) : \Omega \rightarrow 2^{R^m}$  as follows:

$$(3.3) \quad \mathbb{F}(t, x) \equiv \{f(t, x) + B(t, x)u : u \in S(G(t, x) + F(\cdot), h, \varphi, K(\cdot))\},$$

where  $S(G(t, x) + F(\cdot), h, \varphi, K(\cdot))$  denotes the solution set of the generalized mixed quasi-variational inequality in the GDMQVI (1.1). The following lemma presents some properties of the set-valued mapping  $\mathbb{F}$  defined by (3.3) under the hypotheses (A) and (B).

**Lemma 3.4.** Let the hypotheses (A) and (B) hold, the functions  $h, F : R^n \rightarrow R^n$  and the set-valued mapping  $K : R^n \rightrightarrows R^n$  be continuous on  $R^n$ , and for each  $u \in R^n, \varphi : R^n \rightarrow R \cup \{+\infty\}$

be proper and continuous on  $K(u)$ . Suppose that, for all  $q \in G(\Omega)$ ,  $S(q + F(\cdot), h, \varphi, K(\cdot))$  is nonempty and there exists a constant  $\rho > 0$  such that

$$(3.4) \quad \sup\{\|u\| : u \in S(q + F, h, \varphi, K(\cdot))\} \leq \rho(1 + \|q\|).$$

Then there exists a constant  $\rho^{\mathbb{F}} > 0$  such that

$$\sup\{\|y\| : y \in \mathbb{F}(t, x)\} \leq \rho^{\mathbb{F}}(1 + \|x\|), \quad \forall (t, x) \in \Omega.$$

Moreover,  $\mathbb{F}$  is upper semicontinuous and has closed-value on  $\Omega$ .

*Proof.* The proof is very similar to the proof of Lemma 3.1 in [11]. We omit it here. □

**Theorem 3.1.** Let the hypotheses (A) and (B),  $F : R^n \rightarrow R^n$  be Lipschitz continuous with constants  $\rho_1$  on  $R^n$ ,  $h$  be a linear and inverse function with linear coefficient being  $\rho_2$ , and  $\varphi : R^n \rightarrow R \cup \{+\infty\}$  be a proper, convex and continuous functional with  $\varphi(u(\cdot))$  being integral for every integral function  $u \in L^2[0, T]$ . Suppose that

- (i)  $F$  is  $\lambda$ -strongly monotone on  $R^n$  and  $(h, F)$  is  $\mu$ -strongly monotone couple on  $R^n$ ;
- (ii) there exists  $k > 0$  such that  $\|P_{K(u)}z - P_{K(y)}z\| \leq k\|u - y\|, \quad \forall u, y \in R^n, z \in \{v : v = h(u) - p(q + F(u)), u \in R^n, q \in G(\Omega)\}$ .
- (iii)  $(\rho_2^2 - 2p\mu + p^2\rho_1^2)^{1/2} + p(1 - 2\lambda + \rho_1^2)^{1/2} < p - k$ .
- (iv)  $K : R^n \rightrightarrows R^n$  is a continuous set-valued mapping such that, for each  $u \in R^n, K(u) \subset R^n$  is a closed convex set and  $K$  has a linear growth.

Then the initial-value GDMQVI(1.1) has a Carathéodory weak solution.

*Proof.* By the Theorem 4.1 in [12], it follows from the assumptions (i), (ii) and (iii) that  $S(q + F, h, \varphi, K(\cdot))$  is a nonempty singleton for every  $q \in G(\Omega)$ , which is assumed by  $\{u\}$ , and thus  $h(u) \in K(u)$ . Since  $h$  is linear, inverse and  $K$  has a linear growth, there exists a constant  $c > 0$  such that for any  $q \in G(\Omega)$ ,

$$\|u\| = \|h^{-1}h(u)\| \leq c\|h^{-1}\|(1 + \|u\|).$$

Then by Lemma 3.4, the set-valued mapping  $\mathbb{F}$ , defined by (3.3), is an upper semicontinuous set-valued mapping with nonempty closed convex values on  $\Omega$  and there exists a constant  $\rho^{\mathbb{F}} > 0$  such that

$$\sup\{\|y\| : y \in \mathbb{F}(t, x)\} \leq \rho^{\mathbb{F}}(1 + \|x\|), \quad \forall (t, x) \in \Omega.$$

This indicates by Lemma 2.2 that inclusion problem  $DI : \dot{x} \in \mathbb{F}(t, x), x(0) = x_0$  has a weak solution  $x$  in the sense of Carathéodory, which implies that

$$\|x(t)\| \leq \|x_0\| + \int_0^t \rho^{\mathbb{F}}(1 + \|x(s)\|)ds.$$

Moreover, by Gronwall's lemma,

$$\|x(t)\| \leq (\|x_0\| + \rho^{\mathbb{F}}T)e^{\rho^{\mathbb{F}}T}.$$

Now, we prove that  $U(t, x) = S(G(t, x) + F, h, \varphi, K(\cdot))$  is closed on  $\Omega$ . To this end, let  $\{(t_n, x_n)\} \subset \Omega$  be a sequence converging to some vector  $(t_0, x_0) \in \Omega$ , and  $\{u_n\} \subset U(t_n, x_n)$  converging to  $u_0$ . Thus,  $h(u_n) \in K(u_n)$  and

$$(3.5) \quad \langle y - h(u_n), G(t_n, x_n) + F(u_n) \rangle + p\varphi(y) - p\varphi(h(u_n)) \geq 0, \quad \forall y \in K(u_n).$$

Since  $h$  is continuous and  $K$  is upper semi-continuous on  $R^n$ , it follows  $h(u_n) \in K(u_n)$  that  $h(u_0) \in K(u_0)$ . The lower semicontinuity of  $K$  implies that, for any  $\hat{y} \in K(u_0)$ , there exists  $y_n \in K(u_n)$  such that  $y_n \rightarrow \hat{y}$ . This implies by (3.5) that

$$(3.6) \quad \langle y_n - h(u_n), G(t_n, x_n) + F(u_n) \rangle + p\varphi(y_n) - p\varphi(h(u_n)) \geq 0.$$

By letting  $n \rightarrow \infty$  at both sides of the above inequality, we obtain that

$$(3.7) \quad \begin{aligned} & \langle \hat{y} - h(u_0), G(t_0, x_0) + F(u_0) \rangle + p\varphi(\hat{y}) - p\varphi(h(u_0)) \\ & \geq \lim_{n \rightarrow \infty} [\langle y_n - h(u_n), G(t_n, x_n) + F(u_n) \rangle] + p\varphi(y_n) - p \liminf_{n \rightarrow \infty} \varphi(h(u_n)) \\ & \geq 0. \end{aligned}$$

This means that  $u_0 \in U(t_0, x_0)$  and thus  $U(t, x)$  is closed on  $\Omega$ . Then, by Lemma 2.3 we know GDMQVI (1.1) admits a Carathéodory weak solution.  $\square$

#### 4. STABILITY FOR THE GDMQVI

In this section, we aim to study the stability of solutions to the GDMQVI (1.1). For this purpose, we consider the parametric GDMQVI, which is denoted by PGDMQVI, as follows:

$$(4.8) \quad \left\{ \begin{aligned} & \dot{x}(t) = f(t, x(t)) + B(t, x(t))u(t), \\ & \langle y - h(u), G(t, x) + F(u, z) \rangle + p\varphi(y) - p\varphi(h(u)) \geq 0, \quad \forall y \in K(u, z), \\ & h(u) \in K(u, z), \\ & x(0) = x_0, \end{aligned} \right.$$

where, with  $(Z, d)$  being a metric space,  $F : R^n \times Z \rightarrow R^n$  and  $K : R^n \times Z \rightrightarrows R^n$  are the perturbed mappings of the corresponding mappings in the GDMQVI (1.1) respectively. For easy of writing, we denote by  $SD(z)$  the Carathéodory weak solution of the GDMQVI (4.8) and the set of all  $u$  is denoted by  $SD_u(z)$ .

**Theorem 4.2.** *Let the hypotheses (A) and (B) hold,  $h : R^n \rightarrow R^n$  be linear and inverse,  $F : R^n \times Z \rightarrow R^n$  and  $K : R^n \times Z \rightrightarrows R^n$  be continuous mappings such that  $K(\cdot, z)$  have a linear growth for every  $z \in Z$ ,  $\varphi : R^n \rightarrow R \cup \{+\infty\}$  be a proper, convex and continuous functional with  $\varphi(u(t))$  being integral for every  $u \in L^2[0, T]$ , and  $z_0 \in Z$  be a given point. Suppose that*

- (i) *there exists a neighborhood  $U(z_0)$  of  $z_0$  such that, for any  $z \in U(z_0)$  and  $q \in G(\Omega)$ ,  $S(q + F(\cdot, z), h, \varphi, K(\cdot, z))$  is a nonempty singleton;*
- (ii)  *$K$  is closed on  $R^n \times \{z_0\}$  and  $K(R^n \times Z)$  is a bounded set;*
- (iii)  *$SD_u(z)$  is uniformly compact at  $z_0$ .*

Then  $SD(z)$  is upper semicontinuous at  $z_0 \in Z$ .

*Proof.* By Theorem 3.1, it follows from assumptions (i) and (ii) that  $SD(z)$  is nonempty for any  $z \in U(z_0)$ . Now, we prove the assertion. We assume the contrary holds for the sake of contradiction. Then there exists an open set  $N$  containing  $SD(z_0)$  such that, for every sequence  $z_n \in U(z_0)$  with  $z_n \rightarrow z_0$ , there exists  $(x_n, u_n) \in SD(z_n)$  but  $(x_n, u_n) \notin N$ , for every  $n$ . Since  $(x_n, u_n) \in SD(z_n)$ , we get that

- (1) for any  $0 \leq s \leq t \leq T$ ,  $x_n(t) - x_n(s) = \int_s^t [f(\tau, x_n(\tau)) + B(\tau, x_n(\tau))u_n(\tau)]d\tau$ ;
- (2)  $h(u_n(t)) \in K(u_n(t), z_n)$  and for any  $\hat{y} \in K(u_n(t), z_n)$ ,  
 $\langle \hat{y} - h(u_n(t)), G(t, x_n(t)) + F(u_n(t)) \rangle + p\varphi(\hat{y}) - p\varphi(h(u_n(t))) \geq 0$ ;
- (3)  $x_n(0) = x_0$ .

Since  $h$  is linear and inverse and  $K(R^n \times Z)$  is bounded, there exists a constant  $C > 0$  such that, for any  $n$  and  $t \in [0, T]$ ,  $\|u_n(t)\| < C$ . By the hypotheses (A), (B) and the boundedness of  $K(R^n \times Z)$ , it follows from (1) that  $\{x_n\}$  is uniformly bounded with the norm  $\|x\|_1 = \sup_{t \in [0, T]} \|x(t)\|$  and there exists a constant  $M > 0$  such that, for any  $n$ ,

$$\|x_n(t) - x_n(s)\| \leq M|t - s|.$$

Thus, from Arzelá-Ascoli theorem, we know  $\{x_n\}$  has a subsequence, denoted by  $\{x_n\}$ , such that  $x_n \rightarrow \hat{x}$ . Meanwhile, since  $SD_u(z)$  is uniformly compact at  $z_0$ , there exists a

subsequence of  $\{u_n\}$ , denoted by  $\{u_n\}$ , such that  $u_n \rightarrow \hat{u}$ . This means  $(x_n, u_n) \rightarrow (\hat{x}, \hat{u}) \notin N$ . Therefore, for any  $0 \leq s \leq t \leq T$ , (1) and (3) imply that

$$(4.9) \quad \hat{x}(t) - \hat{x}(s) = \int_s^t [f(\tau, \hat{x}(\tau)) + B(\tau, x(\tau))\hat{u}(\tau)]d\tau$$

$$(4.10) \quad \hat{x}(0) = x_0.$$

By the assumption (ii), i.e.,  $K$  is closed on  $R^n \times \{z_0\}$ , we get from (2) that

$$(4.11) \quad h(\hat{u}(t)) \in K(\hat{u}(t), z_0).$$

Moreover, since  $K$  is lower semicontinuous, for any  $y_0 \in K(\hat{u}(t), z_0)$ , there exists a sequence  $\{y_n\} \subset K(u_n(t), z_n)$  such that  $y_n \rightarrow y_0$ . It follows from (2) that, for any  $y_0 \in K(\hat{u}(t), z_0)$ ,

$$(4.12) \quad \begin{aligned} & \langle y_0 - h(\hat{u}(t)), G(t, \hat{x}(t)) + F(\hat{u}(t), z_0) \rangle + p\varphi(y_0) - p\varphi(h(\hat{u}(t))) \\ & \geq \lim_{n \rightarrow \infty} \langle y_n - h(u_n(t)), G(t, x_n(t)) + F(u_n(t), z_n) \rangle + p \liminf_{n \rightarrow \infty} [\varphi(y_n) - p\varphi(h(u_n(t)))] \\ & \geq 0. \end{aligned}$$

It follows (4.9)-(4.12) that  $(\hat{x}, \hat{u}) \in SD(z_0)$ , which contradicts to  $(\hat{x}, \hat{u}) \notin N$ . □

**Theorem 4.3.** *Let the hypotheses (A) and (B) hold for the mappings  $(f, G, B)$ ,  $h = I : R^n \rightarrow R^n$  be the identity function,  $z_0 \in Z$  be a given point,  $F : R^n \times Z \rightarrow R^n$  and  $K : R^n \times Z \rightrightarrows R^n$  be continuous mappings such that  $F(\cdot, z_0)$  is  $\lambda$ -strongly monotone with  $\lambda > 0$  and  $K(\cdot, z)$  have a linear growth for every  $z \in Z$ , and  $\varphi : R^n \rightarrow R \cup \{+\infty\}$  be a proper, convex and continuous functional with  $\varphi(u(t))$  being integral for every  $u \in L^2[0, T]$ . Suppose that*

- (i) *there exists a neighborhood  $U(z_0)$  of  $z_0$  such that  $SD(z)$  is nonempty for any  $z \in U(z_0)$ ;*
- (ii)  *$K$  is closed on  $R^n \times \{z_0\}$ ,  $K(R^n \times Z)$  is bounded, and  $K(\hat{u}(t), z_0) = K(\bar{u}(t), z_0)$  for any  $\hat{u}, \bar{u} \in SD_u(z_0)$ ;*
- (iii)  *$SD_u(z)$  is uniformly compact at  $z_0$ ;*
- (iv) *for any  $(\hat{x}, \hat{u}) \in SD(z_0)$ ,*

$$\langle y - \hat{u}(t), G(t, \hat{x}(t)) + F(\hat{u}(t), z_0) \rangle + p\varphi(y) - p\varphi(\hat{u}(t)) > 0, \quad \forall y \in K(\hat{u}(t), z_0) \setminus \{\hat{u}(t)\}.$$

Then  $SD(z)$  is lower semicontinuous at  $z_0$ .

*Proof.* Suppose, on the contrary, that  $SD(z)$  is not lower semicontinuous at  $z_0$ . Then there exists a sequence  $\{z_n\}$  in  $Z$  with  $z_n \rightarrow z_0$ , and  $(\hat{x}, \hat{u}) \in SD(z_0)$  such that, for every sequence  $(x_n, u_n) \in SD(z_n)$ ,  $(x_n, u_n) \not\rightarrow (\hat{x}, \hat{u})$ . Since  $SD_u(z)$  is uniformly compact at  $z_0$ , there exists a subsequence of  $\{u_n\}$ , denoted by  $\{u_{n_k}\}$ , such that  $u_{n_k} \rightarrow \bar{u}$ . By similar arguments in proof of Theorem 4.2, it is easy to obtain that  $\{x_n\}$  has a subsequence  $\{x_{n_k}\}$  such that  $x_{n_k} \rightarrow \bar{x}$  and  $(\bar{x}, \bar{u}) \in SD(z_0)$ . Thus, for any  $t \in [0, T]$ ,

$$(4.13) \quad \bar{x}(t) = x_0 + \int_0^t [f(\tau, \bar{x}(\tau)) + B(\tau, \bar{x}(\tau))\bar{u}(\tau)]d\tau,$$

$$(4.14) \quad \hat{x}(t) = x_0 + \int_0^t [f(\tau, \hat{x}(\tau)) + B(\tau, \hat{x}(\tau))\hat{u}(\tau)]d\tau.$$

And, by the contradiction assumption,  $(\bar{x}, \bar{u}) \neq (\hat{x}, \hat{u})$ . This means that  $\bar{x} \neq \hat{x}$  or  $\bar{u} \neq \hat{u}$ . Note that if  $\bar{x} \neq \hat{x}$ , it is easy to get from (4.13) and (4.14) that  $\bar{u} \neq \hat{u}$ . If  $\hat{u} \neq \bar{u}$ , it follows from the assumption (iv) that, for all  $y_1 \in K(\hat{u}(t), z_0) \setminus \{\hat{u}(t)\}$ ,

$$(4.15) \quad \langle y_1 - \hat{u}(t), G(t, \hat{x}(t)) + F(\hat{u}(t), z_0) \rangle + p\varphi(y_1) - p\varphi(\hat{u}(t)) > 0$$

and, for all  $y_2 \in K(\bar{u}(t), z_0) \setminus \{\bar{u}(t)\}$ ,

$$(4.16) \quad \langle y_2 - \bar{u}(t), G(t, \bar{x}(t)) + F(\bar{u}(t), z_0) \rangle + p\varphi(y_2) - p\varphi(\bar{u}(t)) > 0.$$

Letting  $y_1 = \bar{u}(t)$  in (4.15) and  $y_2 = \hat{u}(t)$  in (4.16), we get by adding the obtained inequalities that

$$\langle \bar{u}(t) - \hat{u}(t), F(\bar{u}(t), z_0) - F(\hat{u}(t), z_0) \rangle < \langle \bar{u}(t) - \hat{u}(t), G(t, \hat{x}(t)) - G(t, \bar{x}(t)) \rangle.$$

Since  $F : R^n \times Z \rightarrow R^n$  is  $\lambda$ -strongly monotone on  $R^n \times \{z_0\}$  and  $G$  is Lipschitz continuous function on  $\Omega$  with Lipschitz constants  $L_G > 0$ , we get from the above inequality that

$$\|\bar{u}(t) - \hat{u}(t)\| < \frac{L_G}{\lambda} \|\bar{x}(t) - \hat{x}(t)\|.$$

Thus, it follows from (4.13) and (4.14) that there exists a constant  $C_1 > 0$  such that

$$\|\bar{x}(t) - \hat{x}(t)\| < C_1 \int_0^t \|\bar{x}(s) - \hat{x}(s)\| ds.$$

By applying the Gronwall inequality, it is easy to get that  $\|\bar{x}(t) - \hat{x}(t)\| < 0$ , which is a contradiction.  $\square$

**Acknowledgments.** This work was supported by the National Natural Science Foundation of China (11771067, 11501263), State Scholarship Fund of China Scholarship Council (201808510007, 201708515096) Natural Science project of Education Department of Sichuan Province (18ZB0070, 17ZA0030), Youth Science Foundation of Chengdu University of Technology (2017QJ08), China Scholarship Council, Science and Technology Project of Sichuan Province (2017JY0206).

#### REFERENCES

- [1] Agarwal, P., Al-Mdallal, Q., Cho, Y. J. et al., *Fractional differential equations for the generalized Mittag-Leffler function* *Advances in difference equations*, Adv. Differ. Equ., **2018** (2018):58, DOI <https://doi.org/10.1186/s13662-018-1500-7>
- [2] Aubin, J. P. and Frankowska, H., *Set-Valued Analysis*, Birkhäuser, Boston, 1990
- [3] Gwinner, J., *On a new class of differential variational inequalities and a stability result*, Math. Program., **139** (2013), 205–221
- [4] Han, Y., Huang, N. J., Lu, J. et al., *Existence and stability of solutions for inverse variational inequality problems*, Appl. Math. Mech. (English Ed.), **38** (2017), 749–764
- [5] Kaskasem, P., Klin-eam, C., Cho, Y. J., *On the Stability of the Generalized Cauchy-Jensen Set-valued Functional Equations*, J. Fixed Point Theory Appl., (2018) 20: 76, DOI <https://doi.org/10.1007/s11784-018-0558-x>
- [6] Khanh, P. Q., Luu, L. M., *Lower semicontinuity and upper semicontinuity of the solution sets and approximate solution sets of parametric multivalued quasivariational inequalities*, J. Optim. Theory Appl., **133** (2007), 329–339
- [7] Konnov, I. V. and Volotskaya, E. O., *Mixed variational inequalities and economic equilibrium problems*, J. Appl. Math., **2** (2002), 289–314
- [8] Huang, N. J., Lan, H. Y., Cho, Y. J., *Sensitivity analysis for nonlinear generalized mixed implicit equilibrium problems with non-monotone set-valued mappings*, J. Comput. Appl. Math., **196** (2006), 608–618
- [9] Liu, Z. B., Chen, Y. S. et al., *Implicit ishikawa approximation methods for nonexpansive semigroups in CAT(0) spaces*, Abstr. Appl. Anal., **2013**, (2013), doi:10.1155/2013/503198
- [10] Liu, Z. B., Gou, J. H. et al., *A system of generalized variational-hemivariational inequalities with set-valued mappings*, J. Appl. Math., **2013** (2013), 9 pp.
- [11] Li, W., Xiao, Y. B. et al., *A class of differential inverse quasi-variational inequalities in finite dimensional spaces*, J. Nonlinear Sci. Appl., **10** (2017), 4532–4543.
- [12] Li, X. and Zou, Y. Z., *Existence result and error bounds for a new class of inverse mixed quasi-variational inequalities*, J. Inequal. Appl., **1** (2016), 1–13
- [13] Li, X. S., Huang, N. J., O'Regan, D., *Differential mixed variational inequalities in finite dimensional spaces*, Nonlinear Anal., **72** (2010), 3875–3886
- [14] Li, X. S., Huang, N. J. and O'Regan, D., *A class of impulsive differential variational inequalities in finite dimensional spaces*, J. Franklin Inst., **353** (2016), 3151–3175
- [15] Petrusel, A., Petrusel, G. et al., *Fixed point theorems for generalized contractions with applications to coupled fixed point theory*, J. Nonlinear Convex Anal., **19** (2018), 71–87
- [16] Pang, J. S. and Stewart, D., *Differential variational inequalities*, Math. Program. Ser. A, **113** (2008), 345–424
- [17] Pang, J. S., Shen, J., *Strongly regular differential variational systems*, IEEE Trans. Automat. Control, **52** (2007), 242–255

- [18] Stewart, D. E., *Uniqueness for index-one differential variational inequalities*, *Nonlinear Anal. Hybrid Syst.* **2** (2008), 812–818
- [19] Song, Y. S., Muangchoo-in, K. et al., *Successive approximations for common fixed points of a family of  $\alpha$ -nonexpansive mappings*, *J. Fixed Point Theory Appl.*, (2018) **20**: 10. DOI <https://doi.org/10.1007/s11784-018-0483-z>
- [20] Sofonea, M. and Xiao, Y. B., *Fully history-dependent quasivariational inequalities in contact mechanics*, *Appl. Anal.*, **95** (2016), 2464–2484
- [21] Tobin, R. L., *Sensitivity analysis for variational inequalities*, *J. Optim. Theory Appl.*, **48** (1986), 191–209
- [22] Wang, X., Li, W. et al. *Stability for differential mixed variational inequalities*, *Optim. Lett.*, **8** (2013), 1–15
- [23] Wang, X. and Huang, N. J., *Stability analysis for set-valued vector mixed variational inequalities in real reflexive Banach spaces*, *J. Ind. Manag. Optim.*, **9** (2013), 57–74
- [24] Wang, X., Qi, Y. W. et al., *A class of delay differential variational inequalities*, *J. Optim. Theory Appl.*, **172** (2017), 56–69
- [25] Wang, Y. M., Xiao, Y. B. et al., *Equivalence of well-posedness between systems of hemivariational inequalities and inclusion problems*, *J. Nonlinear Sci. Appl.*, **9** (2016), 1178–1192
- [26] Xiao, Y., Fu, X., Zhang, A., *Demand uncertainty and airport capacity choice*, *Transportation Research Part B*, **57** (2013), 91–104
- [27] Xiao, Y. B., Huang, N. J. and Cho, Y. J., *A class of generalized evolution variational inequalities in Banach space*, *Appl. Math. Lett.*, **25** (2012), 914–920
- [28] Xiao, Y. B., Huang, N. J. and Wong, M. M., *Well-posedness of hemivariational inequalities and inclusion problems*, *Taiwanese J. Math.*, **15** (2011), 1261–1276
- [29] Xiao, Y. B., Yang, X. M., Huang and N. J., *Some equivalence results for well-posedness of hemivariational inequalities*, *J. Global Optim.*, **61** (2015), 789–802
- [30] Zhang, W. X., Han, D. R. and Jiang, S. L., *A modified alternating projection based prediction-correction method for structured variational inequalities*, *Appl. Numer. Math.*, **83** (2014), 12–21

<sup>1</sup>DEPARTMENT OF APPLIED MATHEMATICS  
 CHENGDU UNIVERSITY OF TECHNOLOGY  
 CHENGDU, SICHUAN, 610059, CHINA  
 Email address: lovely1w@126.com

<sup>2</sup>UNIVERSITY OF ELECTRONIC SCIENCE AND TECHNOLOGY OF CHINA  
 SCHOOL OF MATHEMATICAL SCIENCES  
 CHENGDU, SICHUAN, 611731 P. R. CHINA  
 Email address: xiaoyb9999@hotmail.com

<sup>3</sup>SCHOOL OF INFORMATION TECHNOLOGY  
 JIANGXI UNIVERSITY OF FINANCE AND ECONOMICS  
 NANCHANG, JIANGXI, 330013, P.R. CHINA  
 Email address: wangxing0793@163.com

<sup>4</sup>CHENGDU UNIVERSITY OF TECHNOLOGY  
 STATE KEY LABORATORY OF GEOHAZARD PREVENTION AND GEOENVIRONMENT PROTECTION  
 CHENGDU, SICHUAN, 610059, CHINA  
 Email address: fengjun@cduet.edu.cn