

Dedicated to Prof. Juan Nieto on the occasion of his 60th anniversary

Convergence results for fixed point iterative algorithms in metric spaces

IOAN A. RUS

ABSTRACT. Let (X, d) be a metric space, $f, f_n : X \rightarrow X$, with $F_f = F_{f_n}$, $n \in \mathbb{N}$. For the fixed point equation

$$(1) \quad x = f(x)$$

we consider the following iterative algorithm,

$$(2) \quad x \in X, x_0 = x, x_{n+1}(x) = f_n(x_n(x)), n \in \mathbb{N}.$$

By definition, the algorithm (2) is convergent if,

$$x_n(x) \rightarrow x^*(x) \in F_f \text{ as } n \rightarrow \infty, \forall x \in X.$$

In this paper we give some conditions on f_n and f which imply the convergence of algorithm (2). In this way we improve some results given in [Rus, I. A., *An abstract point of view on iterative approximation of fixed points: impact on the theory of fixed point equations*, Fixed Point Theory, **13** (2012), No. 1, 179–192]. In our results, in general we do not suppose that, $F_f \neq \emptyset$. Some research directions are formulated.

1. INTRODUCTION

In this paper we study the following two problems:

Problem A. Let (X, d) be a metric space, $f, g : X \rightarrow X$ be such that $F_f = F_g$. For the fixed point equation,

$$(1.1) \quad x = f(x)$$

we consider the following algorithm

$$(1.2) \quad x \in X, x_0 = x, x_{n+1}(x) = g(x_n(x)), n \in \mathbb{N}.$$

By definition, the algorithm (1.2) is convergent if,

$$x_n(x) \rightarrow x^*(x) \in F_f \text{ as } n \rightarrow \infty, \forall x \in X.$$

The convergence of the algorithm (1.2), when f is nonexpansive, X is a bounded, convex and closed subset of a Hilbert, Banach or metric space with a convexity structure ($g(x) = (1 - \lambda)x + \lambda f(x)$, $g(x) = W(x, f(x), \lambda)$, $g(x) = G(x, f(x))$, ...) is studied in an impressive number of papers (see [5], [22], [29], [14], [15], [65], [18], [23], [30], [62], [55], [72], [32], [1], [57], [31], [42], [25], [46], [60], [24], ...).

Problem B. Let (X, d) be a metric space, $f, f_n : X \rightarrow X$, $n \in \mathbb{N}$ be such that, $F_f = F_{f_n}$, $n \in \mathbb{N}$. For the fixed point equation (1.1) we consider the following iterative algorithm,

$$(1.3) \quad x \in X, x_0 = x, x_{n+1}(x) = f_n(x_n(x)), n \in \mathbb{N}.$$

Received: 12.12.2018. In revised form: 20.03.2019. Accepted: 27.03.2019

2010 Mathematics Subject Classification. 47H10, 65J25, 47J25, 65J15, 65J05.

Key words and phrases. *metric space, fixed point, asymptotic regularity, quasicontractive operator, demicompact operator, well posedness of fixed point problem, iterative algorithm, convergence, stability, open problem.*

By definition, the algorithm (1.3) is convergent if,

$$x_n(x) \rightarrow x^*(x) \in F_f \text{ as } n \rightarrow \infty, \forall x \in X.$$

As in the case of Problem A, the convergence of algorithm (1.3), when f is nonexpansive, X is a bounded, convex and closed subset of a Hilbert, Banach or metric space with a convexity structure and f_n are given in the terms of f and the convexity structure of such spaces, is studied in a large number of papers ([28], [5], [29], [22], [3], [8], [10], [12], [13], [16], [17], [23], [30], [37], [39], [40], [41], [43], [66], [64], [26], [36], [20], [1], [34], [67], ...).

In this paper we give some conditions on f and g , respectively on f and f_n which imply the convergence of algorithm (1.2), respectively (1.3). In this way we improve some results given in [54]. In our results, in general we do not suppose apriori that the solution of equation (1.1), F_f , is nonempty. Some research directions are formulated.

2. PRELIMINARIES

2.1. Notations. Throughout this paper we use the same notations as in [54].

2.2. Special classes of sequences in a metric space. Let (X, d) be a metric space. A sequence $(x_n)_{n \in \mathbb{N}}$ in X is called asymptotically regular if,

$$d(x_{n+1}, x_n) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Now, let $f : X \rightarrow X$ be an operator. The sequence $(x_n)_{n \in \mathbb{N}}$ is called f -asymptotically regular if,

$$d(x_n, f(x_n)) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

This two notions are the basic notions in the theory of iterative algorithms (see [12], [5], [22], [34], ...).

2.3. Weakly Picard operators in metric spaces (see [49], [51], [50], [56]). Let (X, d) be a metric space. An operator $f : X \rightarrow X$ is called a weakly Picard operator (*WPO*) if the sequence, $(f^n(x))_{n \in \mathbb{N}}$, converges for all $x \in X$, and its limit, $x^*(x) \in F_f$. If f is *WPO* and, $F_f = \{x^*\}$, then f is called Picard operator (*PO*).

For a *WPO*, $f : X \rightarrow X$, we define the limit operator, $f^\infty : X \rightarrow X$, by $f^\infty(x) = \lim_{n \rightarrow \infty} f^n(x)$. We remark that f^∞ is a retraction on the fixed point set of f , F_f .

An important class of *WPO* is so called, ψ -*WPO*. Let $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be an increasing function, continuous in 0 with $\psi(0) = 0$. The *WPO* f is called ψ -*WPO* iff,

$$d(x, f^\infty(x)) \leq \psi(d(x, f(x))), \forall x \in X.$$

We call a such condition, a retraction-displacement condition.

2.4. Some classes of operators on a metric space. Let (X, d) be a metric space and $f : X \rightarrow X$ be an operator. Then:

- (1) f is an l -contraction if $0 < l < 1$ and

$$d(f(x), f(y)) \leq ld(x, y), \forall x, y \in X;$$

- (2) f is a contractive operator if,

$$d(f(x), f(y)) < d(x, y), \forall x, y \in X, x \neq y;$$

- (3) f is nonexpansive if,

$$d(f(x), f(y)) \leq d(x, y), \forall x, y \in X;$$

- (4) f is Caristi-Browder operator (see [11], [58]) if, f is continuous and there exists, $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$, such that

$$d(x, f(x)) \leq \varphi(x) - \varphi(f(x)), \forall x \in X;$$

- (5) f is quasinonexpansive (see [71], [24], [47]) if $F_f \neq \emptyset$ and

$$d(f(x), x^*) \leq d(x, x^*), \forall x \in X, \forall x^* \in F_f;$$
- (6) f is quasicontractive if $F_f \neq \emptyset$ and

$$d(f(x), x^*) < d(x, x^*), \forall x \in X \setminus F_f, x^* \in F_f;$$
- (7) f is K -demicontractive (see [40], [30], [23], [41], ...) if $K < 1, F_f \neq \emptyset$ and

$$(d(f(x), x^*))^2 \leq (d(x, x^*))^2 + K(d(x, f(x)))^2, \forall x \in X, \forall x^* \in F_f;$$
- (8) f is demicompact (see [45], [35]) if the following implication holds:
 $(x_n)_{n \in \mathbb{N}}$ a bounded sequence in X such that $d(x_n, f(x_n)) \rightarrow 0$ as $n \rightarrow \infty$, implies that there exists a subsequence $(x_{n_i})_{i \in \mathbb{N}}$ which is convergent;
- (9) the fixed point for f is well posed if $F_f = \{x^*\}$ and the following implication holds:
 $(x_n)_{n \in \mathbb{N}}$ in X with $d(x_n, f(x_n)) \rightarrow 0$ as $n \rightarrow \infty$ implies that $x_n \rightarrow x^*$ as $n \rightarrow \infty$;
- (10) the fixed point problem for f is well posed in generalized sense if the following implication holds:
 $(x_n)_{n \in \mathbb{N}}$ a sequence in X such that $d(x_n, f(x_n)) \rightarrow 0$ as $n \rightarrow \infty$, implies that there exists a subsequence $(x_{n_i})_{i \in \mathbb{N}}$, which is convergent to a fixed point of f .

We remark that:

- (a) If f is nonexpansive operator and $F_f = \emptyset$ then f is not a quasinonexpansive operator. In our paper, in what follows, we consider the following notion of quasinonexpansivity. An operator f is quasinonexpansive if or $F_f = \emptyset$ or if $F_f \neq \emptyset$, then

$$d(f(x), x^*) \leq d(x, x^*), \forall x \in X, x^* \in F_f,$$
 i.e., f is quasinonexpansive if

$$d(f(x), x^*) \leq d(x, x^*), \forall x \in X, x^* \in F_f.$$
- (b) If f is continuous and demicompact then the fixed point problem for f is well posed in generalized sense.
- (c) Let (X, d) be a bounded metric space, $f : X \rightarrow X$ be continuous and α_K -condensing operator, where α_K is the Kuratowski measure of noncompactness (see [53], [58], ...). Then the fixed point problem for f is well posed in generalized sense.
- (d) If f is K -demicontractive with $K < 0$, then

$$-K \sum_{n=0}^{\infty} (d(f^n(x), f^{n+1}(x)))^2 \leq (d(x, x^*))^2, \forall x \in X, x^* \in F_f.$$

This condition implies that the sequence $(f^n(x))_{n \in \mathbb{N}}$ is f asymptotically regular, i.e., the operator f is asymptotically regular.

3. DISPLACEMENT CONDITIONS

Let (X, d) be a metric space, $\alpha : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ and $\beta : X \rightarrow \mathbb{R}_+$. By definition, (α, β) is an admissible pair if α satisfies the following implication:

$$(t_n)_{n \in \mathbb{N}} \in \mathbb{R}_+, \alpha(t_n) \rightarrow 0 \Rightarrow t_n \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Now let $g : X \rightarrow X$ be an operator. By definition g satisfies the (α, β) -displacement condition if:

- (1) (α, β) is an admissible pair;
- (2) $\alpha(d(x, g(x))) \leq \beta(x) - \beta(g(x)), \forall x \in X.$

Here are some examples of operators which satisfy a displacement condition:

- (a) If $\alpha(t) = t, \forall t \in \mathbb{R}_+$ and g is continuous then g is a Caristi-Browder operator (see [11], [58]);
- (b) (F.E. Browder [13]) Let $(B, \|\cdot\|)$ be a real Banach space, X be a nonempty, closed, convex subset of B and $g : X \rightarrow X$ be an operator. The following condition appear in [13] on g :

$$\varphi(\|g(x)\|) + \psi(\|x - g(x)\|) \leq \varphi(\|x\|), \forall x \in X,$$

with, $\varphi, \psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ continuous, strict increasing with, $\varphi(0) = \psi(0) = 0$.

A such operator satisfies the (α, β) -displacement condition with, $\alpha(t) = \psi(t), \beta(x) = \varphi(\|x\|)$.

- (c) (Mărușter [40], Hicks-Kubicek [30]) Let $(H, \langle \cdot, \cdot \rangle)$ be a Hilbert space, X be a non-empty subset of H and $g : X \rightarrow X$ be a K -demicontractive operator with $K < 0$. Then g satisfies the (α, β) -displacement condition, with $\alpha(t) = -Kt^2$ and $\beta(x) = \|x - x^*\|^2$.
- (d) In the notion of demicontractivity we suppose that the fixed point set of the operator is nonempty. We can give a type of demicontractivity without this condition in the following way.

Let (M, d) be a metric space, $X \subset M$ be a nonempty subset, $g : X \rightarrow X$ be an operator and $p \in M \setminus X$.

By definition, the operator g is K -demicontractive with respect to the point p , if $K < 1$, and

$$(d(g(x), p))^2 \leq (d(x, p))^2 + K(d(x, g(x)))^2, \forall x \in X.$$

It is clear that if $K < 0$, then g satisfies (α, β) -displacement condition with, $\alpha(t) = -Kt^2$ and $\beta(x) = (d(x, p))^2$.

For a such trick see, for example, [38], [62].

We have the following results in terms of displacement conditions.

Theorem 3.1. *Let (X, d) be a metric space and $g : X \rightarrow X$ be an operator which satisfies the (α, β) -displacement condition. Then the operator g is asymptotically regular.*

Proof. Let $x \in X$. The (α, β) -displacement condition implies that:

$$\begin{aligned} \alpha(d(x, g(x))) &\leq \beta(x) - \beta(g(x)), \\ \alpha(d(g(x), g^2(x))) &\leq \beta(g(x)) - \beta(g^2(x)), \\ &\vdots \\ \alpha(d(g^n(x), g^{n+1}(x))) &\leq \beta(g^n(x)) - \beta(g^{n+1}(x)), \forall n \in \mathbb{N}. \end{aligned}$$

These imply that,

$$\sum_{n=0}^{\infty} \alpha(d(g^n(x), g^{n+1}(x))) \leq \beta(x), \forall x \in X,$$

from which it follows that, g is asymptotically regular. □

Now, let we have two operators, $f, g : X \rightarrow X$ with, $F_f = F_g$. By definition g satisfies (α, β, f) -displacement condition if the pair (α, β) is admissible and

$$\alpha(d(x, f(x))) \leq \beta(x) - \beta(g(x)), \forall x \in X.$$

For a such class of operators we have:

Theorem 3.2. *If g satisfies the (α, β, f) -displacement condition, then the sequence, $(g^n(x))_{n \in \mathbb{N}}$ is f -asymptotically regular.*

Proof. From the (α, β, f) -displacement condition we have that:

$$\begin{aligned} \alpha(d(x, f(x))) &\leq \beta(x) - \beta(g(x)), \forall x \in X, \\ \alpha(d(g(x), f(g(x)))) &\leq \beta(g(x)) - \beta(g^2(x)), \forall x \in X, \\ &\vdots \\ \alpha(d(g^n(x), f(g^n(x)))) &\leq \beta(g^n(x)) - \beta(g^{n+1}(x)), \forall x \in X, \forall n \in \mathbb{N}. \end{aligned}$$

These imply that,

$$\sum_{n=0}^{\infty} \alpha(d(g^n(x), f(g^n(x)))) \leq \beta(x), \forall x \in X.$$

This condition implies that the sequence $(g^n(x))_{n \in \mathbb{N}}$ is f -asymptotically regular for all $x \in X$. □

In the next section we shall use these two results to study the Problem A.

4. THE CONVERGENCE OF THE ALGORITHM IN PROBLEM A

Let (X, d) be a metric space and $f, g : X \rightarrow X$ be two operators with $F_f = F_g$. For the fixed point equation

$$(4.1) \quad x = f(x)$$

we consider the following iterative algorithm:

$$(4.2) \quad x \in X, x_0 = x, x_{n+1}(x) = g(x_n(x)), n \in \mathbb{N}.$$

The problem is in which conditions on f and g the algorithm (4.2) is convergent, i.e., in which conditions on f and g , the operator g is *WPO*?

For a better understanding of the Problem A we start with some examples.

Example 4.1. Let $(B, \|\cdot\|)$ be a Banach space, $X \subset B$ be a nonempty, bounded, closed and convex subset of B and $f : X \rightarrow X$ be a nonexpansive operator. For $\lambda \in]0, 1[$ let f_λ be the Krasnoselski operator, corresponding to f , defined by

$$f_\lambda(x) = (1 - \lambda)x + \lambda f(x).$$

By a Ishikawa Theorem (see [18]) the operator f_λ is asymptotically regular. But,

$$f_\lambda(x) - x = \lambda(f(x) - x), \forall x \in X.$$

This implies that the sequence $(f_\lambda^n(x))_{n \in \mathbb{N}}$ is f -asymptotically regular.

Example 4.2. Let $(B, \|\cdot\|)$ be a Banach space, $f : B \rightarrow B$ be an operator and $\lambda \in \mathbb{R}^* := \mathbb{R} \setminus \{0\}$.

We consider the operator $f_\lambda := (1 - \lambda)1_B + \lambda f$. Then we remark that, $F_f = F_{f_\lambda}$ and f_λ is asymptotically regular if and only if the sequence, $(f_\lambda^n(x))_{n \in \mathbb{N}}$, is f -asymptotically regular.

Example 4.3. Let B be a Banach space and $f : B \rightarrow B$ be an l -Lipschitz operator. In this case, the operator f_λ (see Example 4.2), for $\lambda = \frac{1}{1+l}$ is nonexpansive.

For the Problem A we have:

Theorem 4.1. *We suppose that:*

- (i) g satisfies the (α, β) -displacement condition;
- (ii) there exists $c > 0$ such that,

$$d(x, g(x)) \geq cd(x, f(x)), \forall x \in X;$$

- (iii) the fixed point problem for f is well posed in generalized sense;
- (iv) g is quasinonexpansive.

Then g is WPO and $g^\infty(X) = F_f$.

Proof. Let $x \in X$. From Theorem 3.1, the condition (i) implies that g is asymptotically regular. Condition (ii) implies that the sequence, $(g^n(x))_{n \in \mathbb{N}}$ is f -asymptotically regular. From (iii), there exists a subsequence $(g^{n_i}(x))_{i \in \mathbb{N}}$ such that

$$g^{n_i}(x) \rightarrow x^*(x) \in F_f \text{ as } n \rightarrow \infty.$$

So, $F_f \neq \emptyset$. In this case the condition (iv) is effectively and implies that the sequence $(d(g^n(x), x^*))_{n \in \mathbb{N}}$ is decreasing. So,

$$d(g^n(x), x^*) \rightarrow d \geq 0 \text{ as } n \rightarrow \infty.$$

But, $d(g^{n_i}(x), x^*(x)) \rightarrow 0$ as $n_i \rightarrow \infty$, i.e.,

$$g^n(x) \rightarrow x^*(x) \in F_f \text{ as } n \rightarrow \infty.$$

□

Theorem 4.2. We suppose that:

- (i) g satisfies the (α, β, f) -displacement condition;
- (ii) the fixed point problem for f is well posed in generalized sense;
- (iii) g is quasinonexpansive.

Then g is WPO.

Proof. Let $x \in X$. From Theorem 3.2, condition (i) implies that, $(g^n(x))_{n \in \mathbb{N}}$ is f -asymptotically regular. Now the proof is similar with that of Theorem 4.1. □

In what follows, we give some applications of the above results to the iterative algorithm with admissible perturbation (see [54]).

Following [54] we introduce a new class of operators which generalizes the Krasnosel'ski operators. Let X be a nonempty set, $G : X \times X \rightarrow X$ be an operator. We suppose that:

- (A₁) $G(x, x) = x, \forall x \in X$;
- (A₂) $x, y \in X, G(x, y) = x$ imply, $y = x$.

Let $f : X \rightarrow X$ be an operator. We consider the operator $g = f_G : X \rightarrow X$, defined by

$$f_G(x) := G(x, f(x)).$$

We remark that, $F_f = F_{f_G}$.

We call the operator f_G the admissible perturbation of f corresponding to G . For some examples of admissible perturbation in the case in which X is a subset of linear space, Hilbert space, Banach space, metric space with convexity structure, see [54]. Problem A in this case is the following:

In which conditions on $f : (X, d) \rightarrow (X, d)$ and $G : X \times X \rightarrow X$, the admissible perturbation, f_G of f is WPO?

For some results on this problem in the case of Hilbert and Banach spaces, see: [6], [10], [72], [7], [69], [64], [68], [19], [57], [70], ...

We give a result in a metric space.

Theorem 4.3. We suppose that:

- (i) there exists an admissible pair (α, β) such that:

$$\alpha(d(x, G(x, y))) \leq \beta(x) - \beta(G(x, y)), \forall x, y \in X;$$

(ii) there exists $c > 0$ such that:

$$d(x, G(x, y)) \geq cd(x, y), \forall x, y \in X;$$

(iii) the fixed point problem for f is well posed in generalized sense;

(iv) the operator f_G is quasinonexpansive.

Then, the operator f_G is WPO.

Proof. If, in conditions (i) and (ii) we take $y = f(x)$, then we remark that the operator f_G satisfies the conditions in Theorem 4.1. \square

5. CONVERGENCE OF ALGORITHM IN PROBLEM B

We start with some remarks on the sequences in metric spaces. Let (X, d) be a metric space and $(x_n)_{n \in \mathbb{N}}$ be a sequence in X . We have:

Lemma 5.1. *If there exists an admissible pair, (α, β) , such that,*

$$\alpha(d(x_n, x_{n+1})) \leq \beta(x_n) - \beta(x_{n+1}), \forall n \in \mathbb{N},$$

then the sequence, $(x_n)_{n \in \mathbb{N}}$, is asymptotically regular.

Lemma 5.2. *Let, $f : X \rightarrow X$, be an operator. If there exists an admissible pair, (α, β) , such that,*

$$\alpha(d(x_n, f(x_n))) \leq \beta(x_n) - \beta(x_{n+1}), \forall n \in \mathbb{N},$$

then the sequence, $(x_n)_{n \in \mathbb{N}}$, is f -asymptotically regular.

Now let (X, d) be a metric space and, $f, f_n : X \rightarrow X$ be operators with, $F_f = F_{f_n}$. We consider for the fixed point equation corresponding to f , the algorithm (1.3), i.e.,

$$(5.1) \quad x \in X, x_0 = x, x_{n+1}(x) = f_n(x_n(x)), n \in \mathbb{N}.$$

For this algorithm we have:

Theorem 5.1. *We suppose that:*

(i) there exists an admissible pair such that,

$$\alpha(d(x_n(x), x_{n+1}(x))) \leq \beta(x_n(x)) - \beta(x_{n+1}(x)), \forall n \in \mathbb{N}, \forall x \in X;$$

(ii) $d(x_n(x), x_{n+1}(x)) \geq cd(x_n(x), f(x_n(x)))$, with some $c > 0$, for all $n \in \mathbb{N}$ and $x \in X$;

(iii) the fixed point problem for f is well posed in generalized sense;

(iv) the operators f_n are quasinonexpansive.

Then the sequence, $(x_n(x))_{n \in \mathbb{N}}$ converges to a fixed point of f , $x^(x)$.*

Proof. From (i), the sequence, $x_n(x)$, is asymptotically regular. The condition (ii) implies that, $(x_n(x))_{n \in \mathbb{N}}$ is f -asymptotically regular. Condition, (iii) implies that there exists a subsequence, $(x_{n_i}(x))_{i \in \mathbb{N}}$ of $(x_n(x))_{n \in \mathbb{N}}$ which converges to a fixed point of f , $x^*(x)$. In this case, the condition (iv) is effective and we have that the sequence

$$d(x_n(x), x^*(x)) \rightarrow d \geq 0 \text{ as } n \rightarrow \infty.$$

But, $x_{n_i}(x) \rightarrow x^*(x)$ as $n \rightarrow \infty$. So, the sequence $(x_n(x))_{n \in \mathbb{N}}$ converges to $x^*(x)$. \square

From the above proof it is clear that we have the following result with direct conditions on f and f_n .

Theorem 5.2. *We suppose that:*

(i) f_n satisfies (α, β) -displacement condition, $\forall n \in \mathbb{N}$;

(ii) $d(x, f_n(x)) \geq cd(x, f(x))$, with some $c > 0, \forall x \in X$;

(iii) the fixed point problem for f is well posed in generalized sense;

(iv) f_n is quasinonexpansive, $\forall n \in \mathbb{N}$.

Then, the algorithm (5.1) is convergent.

Proof. From (i) we have that

$$\alpha(d(x, f_n(x))) \leq \beta(x) - \beta(f_n(x)), \forall n \in \mathbb{N}, \forall x \in X.$$

In this relation, instead of x we put, $x_n(x)$, and we have

$$\alpha(d(x_n(x), x_{n+1}(x))) \leq \beta(x_n(x)) - \beta(x_{n+1}(x)), \forall n \in \mathbb{N}, \forall x \in X.$$

From Lemma 5.1, the sequence $(x_n(x))_{n \in \mathbb{N}}$ is asymptotically regular. From (ii), the sequence $(x_n(x))_{n \in \mathbb{N}}$ is f -asymptotically regular. Now, see the proof of Theorem 5.1. \square

In a similar way we have,

Theorem 5.3. We suppose that:

(i) there exists an admissible pair (α, β) such that,

$$\alpha(d(x_n(x), f(x_n(x)))) \leq \beta(x_n(x)) - \beta(x_{n+1}(x)), \forall n \in \mathbb{N}, \forall x \in X;$$

(ii) the fixed point problem for f is well posed in generalized sense;

(iii) f_n is quasinonexpansive, $\forall n \in \mathbb{N}$.

Then, the sequence, $(x_n(x))_{n \in \mathbb{N}}$, converges to a fixed point of f .

Theorem 5.4. We suppose that:

(i) f_n satisfies (α, β, f) -displacement condition, $\forall n \in \mathbb{N}$;

(ii) the fixed point problem for f is well posed in generalized sense;

(iii) f_n is quasinonexpansive, $\forall n \in \mathbb{N}$.

Then, the algorithm (5.1) is convergent.

6. PROBLEMS

From the above considerations the following questions rise:

- 6.1. To construct a theory for K -demicontractive operators with $K < 0$, in a metric space. For the K -demicontractive operators in Hilbert and Banach spaces see: [40], [30], [4], [5], [17], [23], [41], [69], [6], ...
- 6.2. To give new metric conditions which imply asymptotic regularity of an operator, and in general, not convergence of successive approximations. A similar problem in the case of sequences.

Let (X, d) be a metric space, $g : X \rightarrow X$ be an operator and $(x_n)_{n \in \mathbb{N}}$ be a sequence in X . The (α, β) -displacement condition for g (see Theorem 3.1) and (α, β) -displacement condition for $(x_n)_{n \in \mathbb{N}}$ (see Lemma 5.1), imply asymptotic regularity.

The problem is to give other conditions with these properties.

References for asymptotic regularity: [48], [14], [15], [5], [18], [56], [58], [33], [31], [42], [25], [46], [50], [62]. [59], ...

- 6.3. To give, in a metric space, conditions in which asymptotic regularity of an operator (sequence) implies convergence of successive approximations (sequence).

In 1945, J. Dieudonné (see [48]) has given the following result:

Let $f \in C([a, b] \times \mathbb{R}^m, \mathbb{R}^m)$ and the following Cauchy problem corresponding to f :

$$y'(x) = f(x, y(x)), y(a) = y_0.$$

We consider the successive approximations for this problem,

$$y_{n+1}(x) = y_0 + \int_a^x f(s, y_n(s)) ds, n \in \mathbb{N}.$$

If the Cauchy problem has a unique solution, then there exists, $h \in]0, b - a[$ such that the successive approximations sequence converges uniformly to the unique solution of Cauchy problem on $[a, a + h]$, if and only if the sequence $\{y_{n+1} - y_n\}$ converges uniformly to the null function, uniformly on $[a, a + h]$.

In 1976, B.P. Hillam (see [62]) proves the following result:

A function $f \in C([0, 1], [0, 1])$ is weakly Picard function if and only if f is asymptotically regular.

The problem is to give similar results in a metric space.

References: [48], [9], [5], [10], [14], [15], ...

6.4. In which conditions a nonexpansive operator is a graphic contraction ?

One basic problem in the theory of nonexpansive operators is the following:

Let (X, d) be a metric space and $f : X \rightarrow X$ be a nonexpansive operator. In which conditions f is WPO ?

For a better understanding of the relation between nonexpansive operator theory and graphic contraction theory we present the following well known results.

• Theorem of equivalent statements. Let X be a nonempty set and $f : X \rightarrow X$ be an operator. The following statements are equivalent:

- (i) $F_{f^n} = F_f \neq \emptyset$;
- (ii) there exists a metric d on X with respect to which f is WPO;
- (iii) there exists a complete metric on X with respect to which f is a continuous graphic contraction;
- (iv) $F_f \neq \emptyset$ and there exists a metric d on X with respect to which f is asymptotically regular.

• Graphic Contraction Principle. Let (X, d) be a complete metric space, $f : X \rightarrow X$ be an operator and $l \in]0, 1[$. We suppose that:

- (i) $d(f^2(x), f(x)) \leq ld(x, f(x)), \forall x \in X$;
- (ii) f has closed graph.

Then the operator f is WPO.

• Bernstein operators, $B_n : C[0, 1] \rightarrow C[0, 1]$, are nonexpansive and graphic contractions.

The Bernstein operator, $B_n : C[0, 1] \rightarrow C[0, 1]$, is defined by

$$B_n(f)(x) := \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k}, \quad x \in [0, 1].$$

It is well known that, $\|B_n\| = 1$ and

$$\|B_n^2(f) - B_n(f)\| \leq \left(1 - \frac{1}{2^{n-1}}\right) \|f - B_n(f)\|.$$

So, B_n is a graphic contraction and weakly Picard operator.

References: [55], [50], [44], [52], ...

6.5. To study the stability of algorithms in Problem A and B.

For the notion of stability of an iterative algorithm see: [54], [8], [5], [10], [27], [43], [42], [2], [63], [62], ...

REFERENCES

[1] Alber, Y., Reich, S. and Yao, J.-C., *Iterative methods for solving fixed point problems with nonself-mappings in Banach spaces*, Abstract Appl. Anal., 4 (2003), 193–216

- [2] Aoyama, K., Eshita, K. and Takahashi, W., *Iteration processes for nonexpansive mappings in convex metric spaces* In: Proc. Int. Conf. Nonlinear Anal. Convex Anal., Okinawa, 2005, 31–39
- [3] Bachar, M., Dehaishi, B. A. B. and Khamsi, M. A., *Approximate Fixed Points* In: Fixed Point Theory and Graph Theory, 99–138, Elsevier, 2016
- [4] Bauschke, H. H. and Borwein, J. M., *On projection algorithms for solving convex feasibility problems*, SIAM Review, **38** (1996), No. 3, 367–426
- [5] Berinde, V., *Iterative Approximation of Fixed Points*, Springer, 2007
- [6] Berinde, V., *Convergence theorems for fixed point iterative methods defined as admissible perturbations of a nonlinear operator*, Carpathian J. Math., **29** (2013), No. 1, 9–18
- [7] Berinde, V., Khan, A. R. and Păcurar, M., *Convergence theorems for admissible perturbations of pseudocontractive operators*, Miskolc Math. Notes, **15** (2014), No. 2, 27–37
- [8] Berinde, V., Mărușter, Șt. and Rus, I. A., *An abstract point of view on iterative approximation of fixed points of nonself operators*, J. Nonlinear Convex Anal., **15** (2014), No. 5, 851–865
- [9] Berinde, V., Păcurar, M. and Rus, I. A., *From a Dieudonné theorem concerning the Cauchy problem to an open problem in the theory of weakly Picard operators*, Carpathian J. Math., **30** (2014), No. 3, 283–292
- [10] Berinde, V., Petrușel, A., Rus, I. A. and Șerban, M. A., *The retraction-displacement condition in the theory of fixed point equation with a convergent iterative algorithm*, In: Mathematical Analysis, Approximation Theory and Their Applications, Springer, 2016, 75–106
- [11] Berinde, V. and Rus, I. A., *Caristi-Browder operator theory in distance spaces*, In: Fixed Point Theory and Graph Theory, 1–28, Elsevier, 2016
- [12] Borwein, J., Reich, S. and Shafirir, I., *Krasnoselski-Mann iterations in normed spaces*, Canad. Math. Bull., **35** (1992), No. 1, 21–28
- [13] Browder, F. E., *Convergence theorems for sequences of nonlinear operators in Banach spaces*, Math. Zeitschr., **100** (1967), 201–225
- [14] Browder, F. E. and Petryshyn, W. V., *The solution by iteration of nonlinear functional equations in Banach spaces*, Bull. Amer. Math. Soc., **72** (1966), No. 3, 571–575
- [15] Browder, F. E. and Petryshyn, W. V., *Construction of fixed points of nonlinear mappings in Hilbert space*, J. Math. Anal. Appl., **20** (1967), No. 2, 197–228
- [16] Bruck, R. E., *A simple proof of the mean ergodic theorem for nonlinear contractions in Banach spaces*, Israel J. Math., **32** (1979), 107–116
- [17] Bruck, R. E., *Random products of contractions in metric and Banach spaces*, J. Math. Anal. Appl., **88** (1982), 319–332
- [18] Bruck, R. E., *Asymptotic behavior of nonexpansive mappings*, Contemporary Math., **18** (1983), 1–47
- [19] Bunlue, N. and Suantai, S., *Convergence theorems of fixed point iterative methods defined by admissible functions*, Thai J. Math., **13** (2015), No. 3, 527–537
- [20] Ceng, L.-C., Petrușel, A., Yao, J.-C. and Yao, Y., *Hybrid viscosity extragradient method for systems of variational inequalities, fixed point of nonexpansive mappings, zero points of accretive operators in Banach spaces*, Fixed Point Theory, **19** (2018), No. 2, 487–502
- [21] Chaoaha, P. and Chanthorn, P. C., *Fixed point sets through iterationi schemes*, J. Math. Anal. Appl., **386** (2012), 273–277
- [22] Chidume, C., *Geometric Properties of Banach spaces and Nonlinear Iterations*, Springer, 2009
- [23] Chidume, C. E. and Mărușter, Șt., *Iterative methods for the computation of fixed points of demicontractive mappings*, J. Comput. Appl. Math., **234** (2010), 861–882
- [24] Datson, W. G., *Fixed points of quasilinear mappings*, J. Austral. Math. Soc., **13** (1972), 167–170
- [25] Edelstein, M., *A remark on a theorem of M.A. Krasnoselski*, Amer. Math. Monthly, **73** (1966), 509–510
- [26] Eldred, A. A. and Praveen, A., *Convergence of Mann's iteration for relatively nonexpansive mappings*, Fixed Point Theory, **18** (2017), No. 2, 545–554
- [27] Glăvan, V., *Private communication*, 2012
- [28] Goebel, K. and Kirk, W. A., *Topics in Metric Fixed Point Theory*, Cambridge Univ. Press, 1990
- [29] Goebel, K. and Reich, S., *Uniform convexity, Hyperbolic Geometry and Nonexpansive Mapping*, Marcel Dekker, 1984
- [30] Hicks, T. L. and Kubicek, J. D., *On the Mann iteration process in a Hilbert space*, J. Math. Anal. Appl., **59** (1977), 498–504
- [31] Ishikawa, S., *Fixed point and iteration of a non-expansive mapping in a Banach space*, Proc. Amer. Math. Soc., **59** (1976), 65–71
- [32] Kirk, W. A., *Krasnoselskii's iteration process in hyperbolic space*, Num. Funct. Anal. Optimiz., **4** (1981–82), 371–381
- [33] Kirk, W. A., *Approximate fixed points of nonexpansive maps*, Fixed Point Theory, **10** (2009), No. 2, 275–288
- [34] Kohlembach, U., *Some computational aspect of metric fixed point theory*, Nonlinear Anal., **61** (2005), 823–837

- [35] Krichen, B. and O'Regan, D., *On the class of relatively weakly demicompact nonlinear operators*, Fixed Point Theory, **19** (2018), No. 2, 625–630
- [36] Latif, A., Alofi, A. S. M., Al-Mazroofi, A. E. and Yao, J.-C., *General composite iterative methods for general systems of variational inequalities*, Fixed Point Theory, **19** (2018), No. 1, 287–300
- [37] Leuştean, L., *Nonexpansive iterations in uniformly convex W -hyperbolic spaces*, Contemporary Math., **513** (2010), 193–209
- [38] Lin, L.-J. and Takahashi, W., *Attractive point theorems and ergodic theorems for nonlinear mappings in Hilbert spaces*, Taiwanese J. Math., **16** (2012), No. 5, 1763–1779
- [39] Liu Z., Feng, C., Kang, S. M. and Ume, J. S., *Approximating fixed points of nonexpansive mappings in hyperspaces*, Fixed Point Theory and Appl., 2007, ID50596, 9 pp.
- [40] Măruşter, Şt., *The solution by iteration of nonlinear equations in Hilbert spaces*, Proc. Amer. Math. Soc., **63** (1977), No. 1, 69–73
- [41] Măruşter, Şt. and Rus, I. A., *Kannan contractions and strongly demicontractive mappings*, Creative Math. Inf., **24** (2015), No. 2, 171–180
- [42] Ortega, J. M. and Rheinboldt, W. C., *Iterative Solution of Nonlinear Equation in Several Variables*, Acad. Press, New York, 1970
- [43] Petruşel, A. and Rus, I. A., *An abstract point of view on iterative approximation schemes of fixed points for multi-valued operators*, J. Nonlinear Sci. Appl., **6** (2013), 97–107
- [44] Petruşel, A., Rus, I. A. and Şerban, M. A., *Nonexpansive operators as graphic contractions*, J. Nonlinear Convex Anal., **17** (2016), No. 7, 1409–1415
- [45] Petryshyn, W. V., *Construction of fixed points of demicompact mappings in Hilbert space*, J. Math. Anal. Appl., **14** (1966), 276–284
- [46] Petryshyn, W. V. and Williamson, T. E., *Strong and weak convergence of the sequence of successive approximations for quasi-nonexpansive mappings*, J. Math. Anal. Appl., **43** (1973), 459–497
- [47] Roux, D., *Applicazioni quasi non espansive: approssimazione dei punti fissi*, Rendiconti di Matematica, **10** (1977), 597–605
- [48] Rus, I. A., *On a theorem of Dieudonné*, (V. Barbu, Ed.), Diff. Eq. and Control Theory, Longmann, 1991
- [49] Rus, I. A., *Weakly Picard mappings*, Comment. Mat. Univ. Carolinae, **34** (1993), No. 4, 769–773
- [50] Rus, I. A., *Generalized Contractions and Applications*, Cluj Univ. Press, Cluj-Napoca, 2001
- [51] Rus, I. A., *Picard operators and applications*, Sci. Math. Jpn., **58** (2003), 191–219
- [52] Rus, I. A., *Iterates of Bernstein operators, via contraction principle*, J. Math. Anal. Appl., **292** (2004), No. 1, 259–261
- [53] Rus, I. A., *Fixed Point Structure Theory*, Cluj Univ. Press, Cluj-Napoca, 2006
- [54] Rus, I. A., *An abstract point of view on iterative approximation of fixed points: impact on the theory of fixed point equations*, Fixed Point Theory, **13** (2012), No. 1, 179–192
- [55] Rus, I. A., *Properties of the solutions of those equations for which the Krasnoselski iteration converges*, Carpathian J. Math., **28** (2012), No. 2, 329–336
- [56] Rus, I. A., *Relevant clases of weakly Picard operators*, Analele Univ. Vest Timișoara, Mat. Inf., **54** (2016), No. 2, 3–19
- [57] Rus, I. A., *Some problems in the fixed point theory*, Advances in the Theory of Nonlinear Analysis and its Applications, **2** (2018), No. 1, 1–10
- [58] Rus, I. A., Petruşel, A. and Petruşel, G., *Fixed Point Theory*, Cluj Univ. Press, Cluj-Napoca, 2008
- [59] Schott, D., *Basic properties of Fejer monotone sequences*, Rostock Math. Kolloq., **49** (1995), 57–74
- [60] Senter, H. F. and Datson, W. G., *Approximating fixed points of nonexpansive mappings*, Proc. Amer. Math. Soc., **44** (1974), No. 2, 375–380
- [61] Shih, M.-H. and Takahashi, W., *Positive stochastic matrices as contraction maps*, J. Nonlinear Convex Anal., **14** (2013), No. 4, 649–650
- [62] Singh, S. P. and Watson, B., *On convergence results in fixed point theory*, Rend. Sem. Mat. Univ. Politec. Torino, **51** (1993), No. 2, 73–91
- [63] Smale, S., *On the efficiency of algorithms of analysis*, Bull. Amer. Math. Soc., **13** (1985), 87–121
- [64] Şerban, M. A., *Fiber contraction principle with respect to an iterative algorithm*, J. Operators, 2013, ID408791, 6 pp.
- [65] Takahashi, W., *A convexity in metric space and nonexpansive mappings*, Kodai Math. Sem. Rep., **22** (1970), 142–149
- [66] Takahashi, W., *Nonlinear Functional Analysis. Fixed Point Theory and Applications*, Yokohama Publ., Yokohama, 2000
- [67] Thole, R. L., *Iterative techniques for approximation of fixed points of certain nonlinear mappings in Banach spaces*, Pacific J. Math., **53** (1974), 259–266
- [68] Timiş, I., *New stability results of Picard iteration for contractive type mappings*, Fasciculi Math., 2016, Nr. 56, DOI: 10.1515

- [69] Toscano, E. and Vetro, C., *Admissible perturbations of α - ψ -pseudocontractive operators: convergence theorems*, Math. Methods Appl. Sci., **40** (2016), No. 5, 1438–1447
- [70] Toscano, E. and Vetro, C., *Fixed point iterative schemes for variational inequality problems*, J. Convex Anal., **25** (2018), No. 2, 701–715
- [71] Tricomi, F., *Un teorema sulla convergenza delle successioni formate delle successive iterate di una funzione di una variabile reale*, Giorn. Mat. Battoglini, **54** (1916), 1–9
- [72] Țicală, C., *Approximating fixed points of asymptotically demicontractive mapping by iterative schemes defined as admissible perturbations*, Carpathian J. Math., **33** (2017), No. 3, 381–388

DEPARTMENT OF MATHEMATICS

BABEȘ-BOLYAI UNIVERSITY

MIHAIL KOGĂLNICEANU 1, 400084 CLUJ-NAPOCA, ROMANIA

Email address: iarus@math.ubbcluj.ro